Bulletin of the Iranian Mathematical Society
https://doi.org/10.1007/s41980-022-00706-y

ORIGINAL PAPER

Fourier Transforms of Some Special Functions in Terms of Orthogonal Polynomials on the Simplex and Continuous Hahn Polynomials

Esra Güldoğan Lekesiz[1] · Rabia Aktaş[1] · Iván Area[2]

Received: 21 December 2021 / Revised: 18 April 2022 / Accepted: 4 May 2022
© The Author(s) 2022

Abstract
In this paper, Fourier transform of multivariate orthogonal polynomials on the simplex is presented. A new family of multivariate orthogonal functions is obtained using the Parseval's identity and several recurrence relations are derived.

Keywords Jacobi polynomials · Multivariate orthogonal polynomials · Hahn polynomials · Fourier transform · Parseval identity · Hypergeometric function · Recurrence relation

Mathematics Subject Classification 33C50 · 33C70 · 33C45 · 42B10

1 Introduction

The integral transforms are important mathematical tools used in many fields such as vibration analysis, sound engineering, and image processing. They have wide applications in physics, engineering, and in other scientific and mathematical disciplines. The study of orthogonal polynomials and their transformations have been the subject

Communicated by Majid Gazor.

✉ Iván Area
area@uvigo.gal

Esra Güldoğan Lekesiz
esragldgn@gmail.com

Rabia Aktaş
raktas@science.ankara.edu.tr

[1] Faculty of Science, Department of Mathematics, Ankara University, 06100 Tandoğan, Ankara, Turkey

[2] CITMAga, Universidade de Vigo, Departamento de Matemática Aplicada II, E.E. Aeronáutica e do Espazo, Campus As Lagoas-Ourense, 32004 Ourense, Spain

Published online: 10 June 2022

of many papers during the last several years. By the Fourier transform or other integral transforms, some univariate orthogonal polynomials systems which are mapped onto each other exist [1]. As an example, it is known that Hermite functions which are multiplied Hermite polynomials $H_n(x)$ by $\exp\left(-x^2/2\right)$ are eigenfunctions of Fourier transform [2–5]; similarly, Koelink [2] has showed that Jacobi polynomials are mapped onto the continuous Hahn polynomials using the Fourier transform. Also, in [5], it is seen that classical Jacobi polynomials can be mapped onto Wilson polynomials by the Fourier–Jacobi transform. By inspired of Koelink's paper, in [6], Masjed-Jamei et al. have introduced two new families of orthogonal functions using Fourier transforms of the generalized ultraspherical polynomials and the generalized Hermite polynomials. In [7], the Fourier transform of Routh–Romanovski polynomials $J_n^{(u,v)}(x;a,b,c,d)$ has been obtained. Furthermore, in [3, 8], four new examples of finite orthogonal functions have been derived by use of the Fourier transforms of the finite classical orthogonal polynomials $M_n^{(p,q)}(x)$ and $N_n^{(p)}(x)$, and two symmetric sequences of finite orthogonal polynomials and the Parseval's identity (see [9–11] for details of the finite orthogonal polynomials and symmetric sequences of finite orthogonal polynomials). Tratnik [12, 13] presented multivariable generalization both of all continuous and discrete families of the Askey tableau, providing hypergeometric representation, orthogonality weight function which applies with respect to subspaces of lower degree and biorthogonality within a given subspace. In [14], Koelink et al. gave a non-trivial interaction for multivariable continuous Hahn polynomials. Recently, Güldoğan et al. [15] have obtained new families of orthogonal functions by means of Fourier transforms of bivariate orthogonal polynomials introduced by Koornwinder [16] and Fernández et al. [17].

In the present paper, we study the Fourier transformations of the classical polynomials orthogonal by means of the extension of Jacobi weight function to several variables on the simplex T^r. When $r = 1$, the simplex T^r becomes the interval $[0, 1]$ and the corresponding orthogonal polynomials are Jacobi polynomials $P_n^{(\alpha,\beta)}(2x-1)$ on the interval $[0, 1]$. In this case, Koelink [2] has derived the Fourier transform of certain Jacobi polynomials $P_n^{(\alpha,\beta)}(x)$ on the interval $[-1, 1]$ in terms of continuous Hahn polynomials and has discussed some applications. The Fourier transforms of some other bivariate orthogonal polynomials as well as orthogonal polynomials on the triangle have been studied by Güldoğan et al. [15]. By the motivation of these papers, the main aim is to produce a new family of multivariate orthogonal functions.

In [12], Tratnik obtained multivariate orthogonal polynomials on the simplex by taking limit of continuous multivariate Hahn polynomials. As different from the results in Tratnik's paper, in this paper, we handle multivariate orthogonal polynomials on the simplex, and using Fourier transforms of some special orthogonal functions in terms of these polynomials, we get some special orthogonal functions in terms of continuous Hahn polynomials using the method in [2, 15]. While doing these, first, we define specific special functions, so that they are determined with the motivation to use the orthogonality relation of orthogonal polynomials on the simplex in Parseval's identity created with the help of Fourier transform. To give the results on T^r, we shall proceed by induction and we first discuss the results for $r = 1$ and $r = 2$.

The manuscript is organized as follows: in Sect. 2, we introduce some basic definitions and notations. The main results are stated in the third section, where we will obtain Fourier transforms of the multivariate orthogonal polynomials on the r-simplex, write them in terms of continuous Hahn polynomials, define a new family of multivariate orthogonal functions using the Parseval's identity, and derive several recurrence relations for the new family of multivariate orthogonal functions. Finally, the proofs of the main results are given in the fourth section.

2 Basic Definitions and Notations

The univariate Jacobi polynomials $P_n^{(\alpha,\beta)}(x)$ are defined by the explicit representation

$$P_n^{(\alpha,\beta)}(x) = 2^{-n}\sum_{k=0}^{n}\binom{n+\alpha}{k}\binom{n+\beta}{n-k}(x+1)^k(x-1)^{n-k}. \tag{2.1}$$

For $\alpha, \beta > -1$ they are orthogonal on the interval $[-1, 1]$ with respect to the weight function $w(x) = (1-x)^\alpha (1+x)^\beta$ [19, p. 68, Eq. (4.3.2)]

$$\int_{-1}^{1}(1-x)^\alpha (1+x)^\beta P_n^{(\alpha,\beta)}(x) P_m^{(\alpha,\beta)}(x)\,\mathrm{d}x = \frac{2^{\alpha+\beta+1}\Gamma(\alpha+n+1)\Gamma(\beta+n+1)}{n!(\alpha+\beta+2n+1)\Gamma(\alpha+\beta+n+1)}\delta_{m,n},$$

where $m, n \in \mathbb{N}_0 := \mathbb{N}\cup\{0\}$, $\delta_{m,n}$ is the Kronecker delta, and $\Gamma(\lambda)$ is the Gamma function (cf. [20]) defined by

$$\Gamma(\lambda) = \int_0^\infty x^{\lambda-1}e^{-x}\mathrm{d}x, \quad \Re(\lambda) > 0. \tag{2.2}$$

The (p, q) generalized hypergeometric function is defined by (cf. [20])

$${}_pF_q\left(\begin{matrix}a_1, a_2, \ldots, a_p\\ b_1, b_2, \ldots, b_q\end{matrix}\,\middle|\, x\right) = \sum_{k=0}^{\infty}\frac{(a_1)_k (a_2)_k \cdots (a_p)_k}{(b_1)_k (b_2)_k \cdots (b_q)_k}\frac{x^k}{k!}, \tag{2.3}$$

where $(a)_k = \prod_{r=0}^{k-1}(a+r)$, $(a)_0 = 1$, $k \in \mathbb{N}_0$, is the Pochhammer symbol, and the beta function is given by (cf. [20])

$$B(a,b) = \int_0^1 x^{a-1}(1-x)^{b-1}\,\mathrm{d}x = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)}, \qquad \Re(a), \Re(b) > 0. \tag{2.4}$$

The continuous Hahn polynomials are defined in terms of hypergeometric series as [21]

$$p_n(x;a,b,c,d) = i^n \frac{(a+c)_n (a+d)_n}{n!} {}_3F_2\left(\begin{matrix}-n,\ n+a+b+c+d-1,\ a+ix \\ a+c,\ a+d\end{matrix}\Big| 1\right). \quad (2.5)$$

These polynomials can be presented as a limiting case of the Wilson polynomials [21]. Koelink [2] gave a nice survey of the history this family of orthogonal polynomials. Recently, in [22], some applications of continuous Hahn polynomials to quantum mechanics have been presented.

Let $|\boldsymbol{x}| := x_1 + \cdots + x_r$ for $\boldsymbol{x} = (x_1, \ldots, x_r) \in \mathbb{R}^r$. Let V_n^r be the linear space of orthogonal polynomials respect to the weight function $W_{\boldsymbol{\alpha}}(\boldsymbol{x}) = x_1^{\alpha_1} \cdots x_r^{\alpha_r} (1 - |\boldsymbol{x}|)^{\alpha_{r+1}}$, $\alpha = (\alpha_1, \ldots, \alpha_{r+1})$, $\alpha_i > -1$, on the simplex $T^r = \{\boldsymbol{x} \in \mathbb{R}^r : x_1 \geq 0, \ldots, x_r \geq 0, 1 - |\boldsymbol{x}| \geq 0\}$. The elements of the linear space V_n^r satisfy the following partial differential equation (cf. [23]):

$$\sum_{i=1}^{r} x_i (1 - x_i) \frac{\partial^2 P}{\partial x_i^2} - 2 \sum_{1 \leq i < j \leq r} x_i x_j \frac{\partial^2 P}{\partial x_i \partial x_j} + \sum_{i=1}^{r} ((\alpha_i + 1) - (|\boldsymbol{\alpha}| + r + 1) x_i) \frac{\partial P}{\partial x_i} = -n \{n + |\boldsymbol{\alpha}| + r + 1\} P,$$

which has $\binom{n+r-1}{n}$ linear independent polynomial solutions of total degree n. In here, $|\boldsymbol{\alpha}| = \alpha_1 + \cdots + \alpha_{r+1}$. The space V_n^r has several different bases. One of these bases can be introduced as follows. Let $\mathbf{x}_j$ and $\mathbf{n}^j$ be defined as

$$\boldsymbol{x}_0 = 0, \quad \boldsymbol{x}_j = (x_1, \ldots, x_j), \quad |\boldsymbol{x}_j| = x_1 + \cdots + x_j,$$
$$\mathbf{n}^j = (n_j, \ldots, n_r), \quad |\mathbf{n}^j| = n_j + \cdots + n_r, \quad 1 \leq j \leq r,$$

and $\mathbf{n}^{r+1} := 0$.

Also, let $\mathbf{n} = (n_1, \ldots, n_r)$, $|\mathbf{n}| = n_1 + \cdots + n_r = n$, $\boldsymbol{\alpha} = (\alpha_1, \ldots, \alpha_{r+1})$, and $\boldsymbol{\alpha}^j = (\alpha_j, \ldots, \alpha_{r+1})$, $|\boldsymbol{\alpha}^j| = \alpha_j + \cdots + \alpha_{r+1}$, $1 \leq j \leq r+1$. In this case, an orthogonal base of the space V_n^r is of form (cf. [23])

$$P_{\mathbf{n}}^{(\alpha)}(\boldsymbol{x}) = \prod_{j=1}^{r} (1 - |\boldsymbol{x}_{j-1}|)^{n_j} P_{n_j}^{(a_j, b_j)} \left(\frac{2x_j}{1 - |\boldsymbol{x}_{j-1}|} - 1\right), \quad (2.6)$$

where $P_{n_j}^{(a_j,b_j)}(x_j)$ indicates the classical univariate Jacobi polynomial defined in (2.1), $a_j = 2|\mathbf{n}^{j+1}| + |\boldsymbol{\alpha}^{j+1}| + r - j$ and $b_j = \alpha_j$. It follows:

$$\int_{T^r} W_\alpha(\boldsymbol{x})\, P_{\mathbf{n}}^{(\alpha)}(\boldsymbol{x})\, P_{\mathbf{m}}^{(\alpha)}(\boldsymbol{x})\, \mathrm{d}\mathbf{x} = h_{\mathbf{n}}^{(\alpha)} \delta_{\mathbf{n},\mathbf{m}}, \tag{2.7}$$

where $\mathrm{d}\mathbf{x} = \mathrm{d}x_1 \cdots \mathrm{d}x_r$, and

$$h_{\mathbf{n}}^{(\alpha)} = \prod_{j=1}^{r} \frac{\Gamma\left(2\left|\mathbf{n}^{j+1}\right| + \left|\boldsymbol{\alpha}^{j+1}\right| + r - j + 1 + n_j\right)}{n_j!\left(2\left|\mathbf{n}^{j+1}\right| + \left|\boldsymbol{\alpha}^{j+1}\right| + r - j + 1 + \alpha_j + 2n_j\right)} \times \frac{\Gamma\left(1 + \alpha_j + n_j\right)}{\Gamma\left(2\left|\mathbf{n}^{j+1}\right| + \left|\boldsymbol{\alpha}^{j+1}\right| + r - j + 1 + \alpha_j + n_j\right)}. \tag{2.8}$$

When $r = 1$, the simplex T^r becomes the interval $[0, 1]$ and the corresponding orthogonal polynomials in (2.6) are the Jacobi polynomials

$$P_n^{(\alpha)}(x) := P_n^{(\alpha_2,\alpha_1)}(2x - 1).$$

3 Main Results

Our main results are to obtain the Fourier transformation of the classical orthogonal polynomials on the simplex T^r, to define a new family of multivariate orthogonal functions in terms of multivariable Hahn polynomials and, to derive several recurrence relations. Let us consider the function

$$\begin{aligned} &g_r(x_1, \ldots, x_r; n_1, \ldots, n_r, a_1, \ldots, a_{r+1}, \alpha_1, \ldots, \alpha_{r+1}) \\ &\quad = \prod_{j=1}^{r} \left(1 + \tanh x_j\right)^{a_j} \left(1 - \tanh x_j\right)^{|\mathbf{a}^{j+1}|} P_{\mathbf{n}}^{(\alpha)}(\Upsilon_1, \Upsilon_2, \ldots, \Upsilon_r), \end{aligned} \tag{3.1}$$

for $r \geq 1$, where

$$\Upsilon_1(x_1) = \Upsilon_1 = \frac{1 + \tanh x_1}{2},$$
$$\Upsilon_r(x_1, \ldots, x_r) = \Upsilon_r = \frac{(1 - \tanh x_1)(1 - \tanh x_2) \cdots (1 - \tanh x_{r-1})(1 + \tanh x_r)}{2^r},$$

for $r \geq 2$ and

$$\mathbf{n} = (n_1, \ldots, n_r), \ \boldsymbol{\alpha} = (\alpha_1, \ldots, \alpha_{r+1}), \quad \text{and} \quad \left|\mathbf{a}^j\right| = a_j + \cdots + a_{r+1}.$$

For $r = 1$, it gives

$$g_1(x; n, a_1, a_2, \alpha_1, \alpha_2) = (1 + \tanh x)^{a_1} (1 - \tanh x)^{a_2} P_n^{(\alpha_1,\alpha_2)}(\Upsilon_1), \tag{3.2}$$

where a_1, a_2, α_1, and α_2 are real parameters and

$$\Upsilon_1(x) = \Upsilon_1 = \frac{1 + \tanh x}{2}.$$

From (3.1), we can write g_{r+1} in terms of g_r

$$\begin{aligned} & g_{r+1}\left(x_1, \ldots, x_{r+1}; n_1, \ldots, n_{r+1}, a_1, \ldots, a_{r+2}, \alpha_1, \ldots, \alpha_{r+2}\right) \\ & \quad = 2^{-rn_{r+1}} \left(1 + \tanh x_{r+1}\right)^{a_{r+1}} \left(1 - \tanh x_{r+1}\right)^{a_{r+2}} P_{n_{r+1}}^{(\alpha_{r+2}, \alpha_{r+1})}\left(\tanh x_{r+1}\right) \\ & \qquad \times g_r\left(x_1, \ldots, x_r; n_1, \ldots, n_r, a_1, \ldots, a_r, a_{r+1} + a_{r+2} + n_{r+1}, \right. \\ & \qquad \left. \alpha_1, \ldots, \alpha_r, \alpha_{r+1} + \alpha_{r+2} + 2n_{r+1} + 1\right), \end{aligned} \tag{3.3}$$

where $P_n^{(\alpha_1, \alpha_2)}(x)$ stands for classical univariate Jacobi polynomials defined in (2.1).

3.1 The Fourier Transform of *r*-Dimensional Jacobi Polynomials on *r*-Simplex

The Fourier transform for a function $g(x)$ is defined as ([24, p. 111, Eq. (7.1)])

$$\mathcal{F}\left(g\left(x\right)\right) = \int_{-\infty}^{\infty} e^{-i\xi x} g\left(x\right) \mathrm{d}x, \tag{3.4}$$

and the Fourier transform for a function $g(x_1, \ldots, x_r)$ in r variables is defined as ([24, p. 182, Eq. (11.1a)])

$$\mathcal{F}\left(g\left(x_1, \ldots, x_r\right)\right) = \int_{-\infty}^{\infty} \ldots \int_{-\infty}^{\infty} e^{-i\left(\xi_1 x_1 + \cdots + \xi_r x_r\right)} g\left(x_1, \ldots, x_r\right) \mathrm{d}x_1 \ldots \mathrm{d}x_r. \tag{3.5}$$

Theorem 3.1 *The following result holds true:*

$$\begin{aligned} & \mathcal{F}\left(g_{r+1}\left(x_1, \ldots, x_{r+1}; n_1, \ldots, n_{r+1}, a_1, \ldots, a_{r+2}, \alpha_1, \ldots, \alpha_{r+2}\right)\right) \\ & \quad = 2^{a_{r+1} + a_{r+2} - rn_{r+1} - 1} \frac{\left(\alpha_{r+2} + 1\right)_{n_{r+1}}}{n_{r+1}!} B\left(a_{r+1} - \frac{i\xi_{r+1}}{2}, a_{r+2} + \frac{i\xi_{r+1}}{2}\right) \\ & \qquad \times {}_3F_2\left(\left.\begin{matrix} -n_{r+1}, & n_{r+1} + \alpha_{r+1} + \alpha_{r+2} + 1, & a_{r+2} + \frac{i\xi_{r+1}}{2} \\ & \alpha_{r+2} + 1, & a_{r+1} + a_{r+2} \end{matrix} \right| 1\right) \\ & \qquad \times \mathcal{F}\left(g_r\left(x_1, \ldots, x_r; n_1, \ldots, n_r, a_1, \ldots, a_r, a_{r+1} + a_{r+2} + n_{r+1}, \right.\right. \\ & \qquad \left.\left. \alpha_1, \ldots, \alpha_r, \alpha_{r+1} + \alpha_{r+2} + 2n_{r+1} + 1\right)\right). \end{aligned} \tag{3.6}$$

Theorem 3.2 *By substituting Fourier transform of the specific function in the right-hand side of the equality (3.6), we have*

$$\mathcal{F}\left(g_r\left(\boldsymbol{x};\mathbf{n},\mathbf{a},\boldsymbol{\alpha}\right)\right)=2^{r(a_r+a_{r+1}-1)+\sum_{j=1}^{r-1}ja_j}$$
$$\times\prod_{j=1}^{r}\left\{\frac{\left(2\left|\mathbf{n}^{j+1}\right|+\left|\boldsymbol{\alpha}^{j+1}\right|+r-j+1\right)_{n_j}}{n_j!}\Lambda_j^r(\mathbf{a},\boldsymbol{\alpha},\mathbf{n};\xi_j)\right\},\tag{3.7}$$

where

$$\Lambda_j^r(\mathbf{a},\boldsymbol{\alpha},\mathbf{n};\xi_j)=B\left(a_j-\frac{i\xi_j}{2},\ \left|\mathbf{n}^{j+1}\right|+\left|\mathbf{a}^{j+1}\right|+\frac{i\xi_j}{2}\right)$$
$$\times\ {}_3F_2\left(\begin{matrix}-n_j,\ n_j+2\left|\mathbf{n}^{j+1}\right|+\left|\boldsymbol{\alpha}^{j}\right|+r-j+1,\ \left|\mathbf{n}^{j+1}\right|+\left|\mathbf{a}^{j+1}\right|+\frac{i\xi_j}{2}\\ 2\left|\mathbf{n}^{j+1}\right|+\left|\boldsymbol{\alpha}^{j+1}\right|+r-j+1,\ \left|\mathbf{n}^{j+1}\right|+\left|\mathbf{a}^{j}\right|\end{matrix}\Big|\,1\right),$$

or, in terms of the continuous Hahn polynomials (2.5)

$$\Lambda_j^r(\mathbf{a},\boldsymbol{\alpha},\mathbf{n};\xi_j)=\frac{n_j!B\left(a_j-\frac{i\xi_j}{2},\ \left|\mathbf{n}^{j+1}\right|+\left|\mathbf{a}^{j+1}\right|+\frac{i\xi_j}{2}\right)}{i^{n_j}\left(\left|\mathbf{n}^{j+1}\right|+\left|\mathbf{a}^{j}\right|\right)_{n_j}\left(2\left|\mathbf{n}^{j+1}\right|+\left|\alpha^{j+1}\right|+r-j+1\right)_{n_j}}$$
$$\times\ p_{nj}\left(\frac{\xi_j}{2};\left|\mathbf{n}^{j+1}\right|+\left|\mathbf{a}^{j+1}\right|,\alpha_j-a_j+1,\left|\mathbf{n}^{j+1}\right|+\left|\alpha^{j+1}\right|-\left|\mathbf{a}^{j+1}\right|+r-j+1,a_j\right),$$

$g_r\left(\boldsymbol{x};\mathbf{n},\mathbf{a},\boldsymbol{\alpha}\right)$ *is the function given in* (3.1) *for* $r\geq 1$, *and*

$$\boldsymbol{x}=(x_1,\ldots,x_r),\quad \mathbf{n}=(n_1,\ldots,n_r),\quad \mathbf{n}^j=\left(n_j,\ldots,n_r\right),$$
$$\left|\mathbf{n}^j\right|=n_j+\cdots+n_r,\quad \boldsymbol{\alpha}=(\alpha_1,\ldots,\alpha_{r+1}),\quad \boldsymbol{\alpha}^j=\left(\alpha_j,\ldots,\alpha_{r+1}\right),$$
$$\mathbf{a}=(a_1,\ldots,a_{r+1}),\mathbf{a}^j=\left(a_j,\ldots,a_{r+1}\right),\quad \left|\mathbf{a}^j\right|=a_j+\cdots+a_{r+1}.$$

3.2 The Class of Special Functions Using Fourier Transform of *r*-Dimensional Polynomials on the *r*-Simplex

The Parseval's identity corresponding to (3.4) is given by the statement [24, p. 118, Eq. (7.17)]

$$\int_{-\infty}^{\infty}g\left(x\right)\overline{h\left(x\right)}\mathrm{d}x=\frac{1}{2\pi}\int_{-\infty}^{\infty}\mathcal{F}\left(g\left(x\right)\right)\overline{\mathcal{F}\left(h\left(x\right)\right)}\mathrm{d}\xi,$$

and the Parseval's identity corresponding to (3.5) is given by [24, p. 183, (iv)]

$$\int_{-\infty}^{\infty}\cdots\int_{-\infty}^{\infty}g\left(x_1,\ldots,x_r\right)\overline{h\left(x_1,\ldots,x_r\right)}\mathrm{d}x_1\ldots\mathrm{d}x_r$$
$$=\frac{1}{(2\pi)^r}\int_{-\infty}^{\infty}\ldots\int_{-\infty}^{\infty}\mathcal{F}\left(g\left(x_1,\ldots,x_r\right)\right)\overline{\mathcal{F}\left(h\left(x_1,\ldots,x_r\right)\right)}\mathrm{d}\xi_1\cdots\mathrm{d}\xi_r.\tag{3.8}$$

Theorem 3.3 *Let be* $\boldsymbol{n} := (n_1, n_2, \ldots, n_r)$, $\boldsymbol{m} := (m_1, m_2, \ldots, m_r)$, $\left|\mathbf{n}^j\right| = n_j + n_{j+1} + \cdots + n_r$ *and* $\boldsymbol{x} := (x_1, x_2, \ldots, x_r)$ *for* $\boldsymbol{x} \in \mathbb{R}^r$. *In here,* $\mathbf{a} := (a_1, a_2, \ldots, a_{r+1})$ *and* $\mathbf{b} := (b_1, b_2, \ldots, b_{r+1})$. *Let*

$$\begin{aligned} \mathbf{a}^j &= \left(a_j, \ldots, a_{r+1}\right), \quad 1 \le j \le r+1, \\ \mathbf{b}^j &= \left(b_j, \ldots, b_{r+1}\right), \quad 1 \le j \le r+1, \end{aligned}$$

so $\left|\mathbf{a}^j\right| = a_j + a_{j+1} + \cdots + a_{r+1}$ *and* $\left|\mathbf{b}^j\right| = b_j + b_{j+1} + \cdots + b_{r+1}$. *The following equality is satisfied:*

$$\begin{aligned} &\int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} W_r\left(\boldsymbol{x}, \mathbf{a}, \mathbf{b}\right) \, {}_rS_{\mathbf{n}}\left(i\boldsymbol{x}; \mathbf{a}, \mathbf{b}\right) \, {}_rS_{\mathbf{m}}\left(-i\boldsymbol{x}; \mathbf{b}, \mathbf{a}\right) \mathrm{d}\mathbf{x} \\ &\quad = 2^{2r} \pi^r h_{\mathbf{n}}^{(\mathbf{a}+\mathbf{b}-\mathbf{1})} \prod_{j=1}^{r} \frac{\left(n_j!\right)^2 \Gamma\left(\left|\mathbf{n}^{j+1}\right| + \left|\mathbf{a}^j\right|\right) \Gamma\left(\left|\mathbf{n}^{j+1}\right| + \left|\mathbf{b}^j\right|\right)}{\left(2\left|\mathbf{n}^{j+1}\right| + \left|\mathbf{a}^{j+1}\right| + \left|\mathbf{b}^{j+1}\right|\right)_{n_j}^2} \delta_{n_j, m_j}, \end{aligned}$$

where

$$\begin{aligned} &W_r\left(\boldsymbol{x}, \mathbf{a}, \mathbf{b}\right) := W_r\left(x_1, \ldots, x_r; a_1, \ldots, a_{r+1}, b_1, \ldots, b_{r+1}\right) \\ &\quad = \prod_{j=1}^{r} \left\{ \Gamma\left(a_j - \frac{ix_j}{2}\right) \Gamma\left(\left|\mathbf{a}^{j+1}\right| + \frac{ix_j}{2}\right) \Gamma\left(b_j + \frac{ix_j}{2}\right) \Gamma\left(\left|\mathbf{b}^{j+1}\right| - \frac{ix_j}{2}\right) \right\} \end{aligned}$$

for $a_j, b_j > 0$ *;* $j = 1, 2, \ldots, r+1$ *and*

$$\begin{aligned} {}_rS_{\mathbf{n}}\left(\boldsymbol{x}; \mathbf{a}, \mathbf{b}\right) &= \prod_{j=1}^{r} \left\{ \left(\left|\mathbf{a}^{j+1}\right| + \frac{x_j}{2}\right)_{\left|\mathbf{n}^{j+1}\right|} \right. \\ &\quad \left. \times {}_3F_2\left(\begin{matrix} -n_j, n_j + 2\left|\mathbf{n}^{j+1}\right| + \left|\mathbf{a}^j\right| + \left|\mathbf{b}^j\right| - 1, \left|\mathbf{n}^{j+1}\right| + \left|\mathbf{a}^{j+1}\right| + \frac{x_j}{2} \\ 2\left|\mathbf{n}^{j+1}\right| + \left|\mathbf{a}^{j+1}\right| + \left|\mathbf{b}^{j+1}\right|, \left|\mathbf{n}^{j+1}\right| + \left|\mathbf{a}^j\right| \end{matrix} \Big| 1 \right) \right\} \end{aligned} \tag{3.9}$$

or, in terms of Hahn polynomials (2.5)

$$\begin{aligned} {}_rS_{\mathbf{n}}\left(\boldsymbol{x}; \mathbf{a}, \mathbf{b}\right) &= \prod_{j=1}^{r} \left\{ \frac{n_j! i^{-n_j}}{\left(\left|\mathbf{n}^{j+1}\right| + \left|\mathbf{a}^j\right|\right)_{n_j} \left(2\left|\mathbf{n}^{j+1}\right| + \left|\mathbf{a}^{j+1}\right| + \left|\mathbf{b}^{j+1}\right|\right)_{n_j}} \right. \\ &\quad \left. \times \left(\left|\mathbf{a}^{j+1}\right| + \frac{x_j}{2}\right)_{\left|\mathbf{n}^{j+1}\right|} p_{n_j}\left(\frac{-ix_j}{2}; \left|\mathbf{n}^{j+1}\right| + \left|\mathbf{a}^{j+1}\right|, b_j, \left|\mathbf{n}^{j+1}\right| + \left|\mathbf{b}^{j+1}\right|, a_j\right) \right\} \end{aligned}$$

for $r \ge 1$.

Remark 3.4 The weight function of this orthogonality relation is positive when all equalities hold simultaneously: $a_j = b_j$ for $j = 1, 2, \ldots, r+1$.

3.3 Recurrence Relations for the Functions ${}_rS_{\mathbf{n}}(\mathbf{x}; \mathbf{a}, \mathbf{b})$

We now derive recurrence relations for the special functions ${}_rS_{\mathbf{n}}(\mathbf{x}; \mathbf{a}, \mathbf{b})$. First, we recall the following relations for hypergeometric function ${}_3F_2\left(\begin{matrix} m_1, m_2, m_3 \\ s_1, s_2 \end{matrix} \middle| z\right)$ that might be obtained by considering the Zeilberger's algorithm [25] based on the Fasenmyer seminal works [26].

Lemma 3.5 *For $|z| < 1$, the hypergeometric function ${}_3F_2$ satisfies the following known recurrence relations:*

(i)

$$\begin{aligned}(z-1)\ {}_3F_2\left(\begin{matrix} m_1+1, m_2, m_3 \\ s_1, s_2 \end{matrix} \middle| z\right) &= (B_1 + C_1 z)\ {}_3F_2\left(\begin{matrix} m_1, m_2, m_3 \\ s_1, s_2 \end{matrix} \middle| z\right) \\ &+ (B_2 + C_2 z)\ {}_3F_2\left(\begin{matrix} m_1-1, m_2, m_3 \\ s_1, s_2 \end{matrix} \middle| z\right) + B_3\, {}_3F_2\left(\begin{matrix} m_1-2, m_2, m_3 \\ s_1, s_2 \end{matrix} \middle| z\right),\end{aligned} \tag{3.10}$$

where

$$\begin{aligned}B_1 &= \frac{s_1+s_2+1-3m_1}{m_1},\ B_2 = \frac{s_1 s_2 + (m_1-1)\,(3m_1 - 2\,(s_1+s_2+1))}{(m_1-1)_2}, \\ B_3 &= -\frac{(m_1-s_1-1)\,(m_1-s_2-1)}{(m_1-1)_2}, \\ C_1 &= \frac{2m_1-m_2-m_3-1}{m_1},\ C_2 = -\frac{(m_1-1)\,(m_1-m_2-m_3-1)+m_2 m_3}{(m_1-1)_2};\end{aligned}$$

(ii)

$$\begin{aligned}(z-1)\ {}_3F_2\left(\begin{matrix} m_1, m_2, m_3 \\ s_1-1, s_2 \end{matrix} \middle| z\right) &= (B_1 + C_1 z)\ {}_3F_2\left(\begin{matrix} m_1, m_2, m_3 \\ s_1, s_2 \end{matrix} \middle| z\right) \\ &+ (B_2 + C_2 z)\ {}_3F_2\left(\begin{matrix} m_1, m_2, m_3 \\ s_1+1, s_2 \end{matrix} \middle| z\right) + C_3 z\, {}_3F_2\left(\begin{matrix} m_1, m_2, m_3 \\ s_1+2, s_2 \end{matrix} \middle| z\right),\end{aligned} \tag{3.11}$$

where

$$\begin{aligned}B_1 &= \frac{s_2-2s_1}{s_1-1},\ B_2 = \frac{s_1-s_2+1}{s_1-1},\ C_1 = \frac{3s_1-m_1-m_2-m_3}{s_1-1}, \\ C_3 &= \frac{(s_1-m_1+1)\,(s_1-m_2+1)\,(s_1-m_3+1)}{(s_1-1)_3},\end{aligned}$$

$$C_2 = \frac{(2s_1+1)(m_1+m_2+m_3) - 3(s_1-1)(s_1+2)}{(s_1-1)_2} - \frac{7+m_1m_2+m_1m_3+m_2m_3}{(s_1-1)_2};$$

(iii)

$$s_1\, {}_3F_2\left(\begin{matrix}m_1, m_2, m_3\\ s_1, s_2\end{matrix}\,\middle|\, z\right) + (m_1 - s_1)\, {}_3F_2\left(\begin{matrix}m_1, m_2, m_3\\ s_1+1, s_2\end{matrix}\,\middle|\, z\right) = m_1\, {}_3F_2\left(\begin{matrix}m_1+1, m_2, m_3\\ s_1+1, s_2\end{matrix}\,\middle|\, z\right). \tag{3.12}$$

To derive recurrence relations, by taking into account (3.9), we first give the following relationships between special functions ${}_rS_{\mathbf{n}}(\mathbf{x};\mathbf{a},\mathbf{b})$ as follows:

$$\begin{aligned}&{}_rS_{\mathbf{n}}(\mathbf{x};\mathbf{a},\mathbf{b})\\ &= {}_3F_2\left(\begin{matrix}-n_r,\ n_r+\left|\mathbf{a}^r\right|+\left|\mathbf{b}^r\right|-1,\ a_{r+1}+\frac{x_r}{2}\\ a_{r+1}+b_{r+1},\ \left|\mathbf{a}^r\right|\end{matrix}\,\middle|\, 1\right)\prod_{j=1}^{r-1}\left(\left|\mathbf{a}^{j+1}\right|+\frac{x_j}{2}\right)_{n_r}\\ &\quad\times_{r-1}S_{n_1,\ldots,n_{r-1}}(x_1,\ldots,x_{r-1};a_1,\ldots,a_{r-1},a_r+a_{r+1}+n_r,\\ &\quad b_1,\ldots,b_{r-1},b_r+b_{r+1}+n_r)\end{aligned} \tag{3.13}$$

or

$$\begin{aligned}{}_rS_{\mathbf{n}}(\mathbf{x};\mathbf{a},\mathbf{b}) &= \left(\left|\mathbf{a}^2\right|+\frac{x_1}{2}\right)_{|\mathbf{n}^2|}\ {}_{r-1}S_{n_2,\ldots,n_r}\left(x_2,\ldots,x_r;\mathbf{a}^2,\mathbf{b}^2\right)\\ &\quad\times\ {}_3F_2\left(\begin{matrix}-n_1,\ n_1+2\left|\mathbf{n}^2\right|+|\mathbf{a}|+|\mathbf{b}|-1,\ \left|\mathbf{n}^2\right|+\left|\mathbf{a}^2\right|+\frac{x_1}{2}\\ 2\left|\mathbf{n}^2\right|+\left|\mathbf{a}^2\right|+\left|\mathbf{b}^2\right|,\ \left|\mathbf{n}^2\right|+|\mathbf{a}|\end{matrix}\,\middle|\, 1\right),\end{aligned} \tag{3.14}$$

for $r \geq 2$ and

$${}_1S_{n_1}(x_1;a_1,a_2,b_1,b_2) = {}_3F_2\left(\begin{matrix}-n_1,\ a_2+x_1/2,\ n_1+a_1+a_2+b_1+b_2-1\\ a_2+b_2,\ a_1+a_2\end{matrix}\,\middle|\, 1\right)$$

for $r = 1$.

Proposition 3.6 *For $r \geq 1$, the family of the special function ${}_rS_{\mathbf{n}}(\mathbf{x};\mathbf{a},\mathbf{b})$ satisfies the following recurrence relation:*

$$\begin{aligned}&(B_1+C_1)\ {}_rS_{\mathbf{n}}(\mathbf{x};\mathbf{a},\mathbf{b}) + (B_2+C_2)\ {}_rS_{n_1+1,n_2,\ldots,n_r}(\mathbf{x};\mathbf{a},b_1-1,b_2,\ldots,b_{r+1})\\ &\quad + B_3\ {}_rS_{n_1+2,n_2,\ldots,n_r}(\mathbf{x};\mathbf{a},b_1-2,b_2,\ldots,b_{r+1}) = 0,\end{aligned}$$

where

$$\begin{aligned}
B_1 + C_1 &= \frac{b_1 - 1 + \frac{x_1}{2}}{n_1}, \\
B_3 &= -\frac{\left(n_1 + 2\left|\mathbf{n}^2\right| + \left|\mathbf{a}^2\right| + \left|\mathbf{b}^2\right| + 1\right)(|\mathbf{n}| + |\mathbf{a}| + 1)}{(n_1)_2}, \\
B_2 + C_2 &= \frac{\left(2\left|\mathbf{n}^2\right| + \left|\mathbf{a}^2\right| + \left|\mathbf{b}^2\right|\right)\left(\left|\mathbf{n}^2\right| + |\mathbf{a}|\right)}{(n_1)_2} \\
&\quad - \frac{\left(n_1 + 2\left|\mathbf{n}^2\right| + |\mathbf{a}| + |\mathbf{b}| - 1\right)\left(\left|\mathbf{n}^2\right| + \left|\mathbf{a}^2\right| + \frac{x_1}{2}\right)}{(n_1)_2} \\
&\quad + \frac{n_1 + 3\left|\mathbf{n}^2\right| + a_1 + 2\left|\mathbf{a}^2\right| - b_1 + \left|\mathbf{b}^2\right| + 2 - \frac{x_1}{2}}{n_1}.
\end{aligned}$$

Proposition 3.7 *For $r \geq 1$, the family of the special function ${}_rS_{\mathbf{n}}(\mathbf{x}; \mathbf{a}, \mathbf{b})$ satisfies the following recurrence relation:*

$$\begin{aligned}
&(B_1 + C_1)\ {}_rS_{\mathbf{n}}(\mathbf{x}; \mathbf{a}, \mathbf{b}) + (B_2 + C_2)\ {}_rS_{\mathbf{n}}(\mathbf{x}; \mathbf{a}, b_1 - 1, b_2, \ldots, b_{r+1}) \\
&\quad + B_3\ {}_rS_{\mathbf{n}}(\mathbf{x}; \mathbf{a}, b_1 - 2, b_2, \ldots, b_{r+1}) = 0,
\end{aligned}$$

where

$$\begin{aligned}
B_1 + C_1 &= -\frac{b_1 - 1 + \frac{x_1}{2}}{n_1 + 2\left|\mathbf{n}^2\right| + |\mathbf{a}| + |\mathbf{b}| - 1}, \\
B_3 &= -\frac{(n_1 + a_1 + b_1 - 2)(|\mathbf{n}| + |\mathbf{b}| - 2)}{\left(n_1 + 2\left|\mathbf{n}^2\right| + |\mathbf{a}| + |\mathbf{b}| - 2\right)_2}, \\
B_2 + C_2 &= \frac{\left(2\left|\mathbf{n}^2\right| + \left|\mathbf{a}^2\right| + \left|\mathbf{b}^2\right|\right)\left(\left|\mathbf{n}^2\right| + |\mathbf{a}|\right) + n_1\left(\left|\mathbf{n}^2\right| + \left|\mathbf{a}^2\right| + \frac{x_1}{2}\right)}{\left(n_1 + 2\left|\mathbf{n}^2\right| + |\mathbf{a}| + |\mathbf{b}| - 2\right)_2} \\
&\quad + \frac{n_1 - \left|\mathbf{n}^2\right| - \left|\mathbf{a}^2\right| + 2b_1 - 3 + \frac{x_1}{2}}{n_1 + 2\left|\mathbf{n}^2\right| + |\mathbf{a}| + |\mathbf{b}| - 1}.
\end{aligned}$$

Proposition 3.8 *For the family of the special function ${}_rS_{\mathbf{n}}(\mathbf{x}; \mathbf{a}, \mathbf{b})$, we get*

$$\begin{aligned}
&(B_2 + C_2)\ {}_rS_{\mathbf{n}}(\mathbf{x}; a_1, \ldots, a_{r-1}, a_r + 1, a_{r+1} - 1, b_1, \ldots, b_{r-1}, b_r - 1, b_{r+1} + 1) \\
&\quad + B_3\ {}_rS_{\mathbf{n}}(\mathbf{x}; a_1, \ldots, a_{r-1}, a_r + 2, a_{r+1} - 2, b_1, \ldots, b_{r-1}, b_r - 2, b_{r+1} + 2) \\
&\quad + (B_1 + C_1)\ {}_rS_{\mathbf{n}}(\mathbf{x}; \mathbf{a}, \mathbf{b}) = 0,
\end{aligned}$$

for $r \geq 1$ where

$$\begin{aligned}
B_1 + C_1 &= \left(b_r - 1 + \frac{x_r}{2}\right)\left(a_{r+1} + \frac{x_r}{2} - 1\right), \\
B_3 &= \left(a_r - \frac{x_r}{2} + 1\right)\left(b_{r+1} - \frac{x_r}{2} + 1\right),
\end{aligned}$$

$$B_2 + C_2 = \left(\left|\mathbf{a}^r\right| - b_r + b_{r+1} + 2 - x_r\right)\left(a_{r+1} + \frac{x_r}{2} - 1\right) - \left|\mathbf{a}^r\right|\left(a_{r+1} + b_{r+1}\right) - n_r\left(n_r + \left|\mathbf{a}^r\right| + \left|\mathbf{b}^r\right| - 1\right).$$

Proposition 3.9 *For the family of the special function* ${}_rS_{\mathbf{n}}(\mathbf{x};\mathbf{a},\mathbf{b})$

$$\begin{aligned}&(B_1 + C_1)\ {}_rS_{\mathbf{n}}(\mathbf{x};\mathbf{a},\mathbf{b})\\ &+ (B_2 + C_2)\ {}_rS_{\mathbf{n}}(\mathbf{x};a_1+1,a_2,\ldots,a_{r+1},b_1-1,b_2,\ldots,b_{r+1})\\ &+ C_3\ {}_rS_{\mathbf{n}}(\mathbf{x};a_1+2,a_2,\ldots,a_{r+1},b_1-2,b_2,\ldots,b_{r+1}) = 0,\end{aligned}$$

holds for $r \geq 1$ *where*

$$\begin{aligned}B_1 + C_1 &= -\frac{b_1 - 1 + \frac{x_1}{2}}{\left|\mathbf{n}^2\right| + |\mathbf{a}| - 1},\\ C_3 &= -\frac{(|\mathbf{n}| + |\mathbf{a}| + 1)\,(|\mathbf{n}| + |\mathbf{b}| - 2)\left(a_1 - \frac{x_1}{2} + 1\right)}{\left(\left|\mathbf{n}^2\right| + |\mathbf{a}| - 1\right)_3},\\ B_2 + C_2 &= \frac{\left(2\left|\mathbf{n}^2\right| + 2|\mathbf{a}| + 1\right)\left(2\left|\mathbf{n}^2\right| + \left|\mathbf{a}^2\right| + |\mathbf{b}| - 2 + \frac{x_1}{2}\right)}{\left(\left|\mathbf{n}^2\right| + |\mathbf{a}| - 1\right)_2}\\ &\quad + \frac{n_1\left(n_1 + 2\left|\mathbf{n}^2\right| + |\mathbf{a}| + |\mathbf{b}| - 1\right)}{\left(\left|\mathbf{n}^2\right| + |\mathbf{a}| - 1\right)_2}\\ &\quad - \frac{\left(2\left|\mathbf{n}^2\right| + |\mathbf{a}| + |\mathbf{b}| - 1\right)\left(\left|\mathbf{n}^2\right| + \left|\mathbf{a}^2\right| + \frac{x_1}{2}\right)}{\left(\left|\mathbf{n}^2\right| + |\mathbf{a}| - 1\right)_2}\\ &\quad - \frac{2\left|\mathbf{n}^2\right| + \left|\mathbf{a}^2\right| + \left|\mathbf{b}^2\right| - 1}{\left|\mathbf{n}^2\right| + |\mathbf{a}| - 1}.\end{aligned}$$

Proposition 3.10 *For the family of the special function* ${}_rS_{\mathbf{n}}(\mathbf{x};\mathbf{a},\mathbf{b})$*, we have*

$$\begin{aligned}&(B_1 + C_1)\ {}_rS_{\mathbf{n}}(\mathbf{x};\mathbf{a},\mathbf{b}) + (B_2 + C_2)\ {}_rS_{\mathbf{n}}(\mathbf{x};\mathbf{a},b_1,\ldots,b_{r-1},b_r-1,b_{r+1}+1)\\ &+ C_3\ {}_rS_{\mathbf{n}}(\mathbf{x};\mathbf{a},b_1,\ldots,b_{r-1},b_r-2,b_{r+1}+2) = 0,\end{aligned}$$

for $r \geq 1$ *where*

$$\begin{aligned}B_1 + C_1 &= -\frac{b_r - 1 + \frac{x_r}{2}}{a_{r+1} + b_{r+1} - 1},\\ C_3 &= -\frac{(n_r + a_r + b_r - 2)\,(n_r + a_{r+1} + b_{r+1} + 1)\left(b_{r+1} - \frac{x_r}{2} + 1\right)}{(a_{r+1} + b_{r+1} - 1)_3},\\ B_2 + C_2 &= \frac{n_r\,(n_r + |\mathbf{a}^r| + |\mathbf{b}^r| - 1) - \left(a_{r+1} + \frac{x_r}{2}\right)(|\mathbf{a}^r| + |\mathbf{b}^r| - 1)}{(a_{r+1} + b_{r+1} - 1)_2}\\ &\quad + \frac{(2a_{r+1} + 2b_{r+1} + 1)\left(|\mathbf{a}^r| + b_r - 2 + \frac{x_r}{2}\right)}{(a_{r+1} + b_{r+1} - 1)_2} - \frac{|\mathbf{a}^r| - 1}{a_{r+1} + b_{r+1} - 1}.\end{aligned}$$

Proposition 3.11 *The family of the special function* ${}_rS_{\mathbf{n}}(\mathbf{x};\mathbf{a},\mathbf{b})$ *satisfies the following recurrence relation:*

$$\begin{aligned}
&\left(\left|\mathbf{n}^2\right|+|\mathbf{a}|\right)\ {}_rS_{\mathbf{n}}(\mathbf{x};\mathbf{a},\mathbf{b})\\
&-\left(n_1+2\left|\mathbf{n}^2\right|+|\mathbf{a}|+|\mathbf{b}|-1\right)\ {}_rS_{\mathbf{n}}(\mathbf{x};a_1+1,a_2,\ldots,a_{r+1},b_1,\ldots,b_{r+1})\\
&+(|\mathbf{n}|+|\mathbf{b}|-1)\ {}_rS_{\mathbf{n}}(\mathbf{x};a_1+1,a_2,\ldots,a_{r+1},b_1-1,b_2,\ldots,b_{r+1})=0
\end{aligned}$$

for $r\geq 1$.

4 Proof of Theorems and Propositions

4.1 Proof of Theorem 3.1

Using the equation (3.3) and the equation in [19, p. 62, Eq. (4.21.2)], the Fourier transform of the function g_{r+1} is as follows:

$$\begin{aligned}
&\mathcal{F}(g_{r+1}(x_1,\ldots,x_{r+1};n_1,\ldots,n_{r+1},a_1,\ldots,a_{r+2},\alpha_1,\ldots,\alpha_{r+2}))\\
&=\mathcal{F}\Big(2^{-rn_{r+1}}(1+\tanh x_{r+1})^{a_{r+1}}(1-\tanh x_{r+1})^{a_{r+2}}P_{n_{r+1}}^{(\alpha_{r+2},\alpha_{r+1})}(\tanh x_{r+1})\\
&\quad\times g_r(x_1,\ldots,x_r;n_1,\ldots,n_r,a_1,\ldots,a_r,a_{r+1}+a_{r+2}+n_{r+1},\\
&\quad\alpha_1,\ldots,\alpha_r,\alpha_{r+1}+\alpha_{r+2}+2n_{r+1}+1))\\
&=2^{-rn_{r+1}}\int_{-\infty}^{\infty}\cdots\int_{-\infty}^{\infty}e^{-i(\xi_1x_1+\cdots+\xi_rx_r+\xi_{r+1}x_{r+1})}(1+\tanh x_{r+1})^{a_{r+1}}\\
&\quad\times(1-\tanh x_{r+1})^{a_{r+2}}P_{n_{r+1}}^{(\alpha_{r+2},\alpha_{r+1})}(\tanh x_{r+1})\\
&\quad\times g_r(x_1,\ldots,x_r;n_1,\ldots,n_r,a_1,\ldots,a_r,a_{r+1}+a_{r+2}+n_{r+1},\\
&\quad\alpha_1,\ldots,\alpha_r,\alpha_{r+1}+\alpha_{r+2}+2n_{r+1}+1)\,\mathrm{d}x_{r+1}\mathrm{d}x_r\cdots\mathrm{d}x_1\\
&=2^{-rn_{r+1}}\mathcal{F}(g_r(x_1,\ldots,x_r;n_1,\ldots,n_r,a_1,\ldots,a_r,a_{r+1}+a_{r+2}+n_{r+1},\\
&\quad\alpha_1,\ldots,\alpha_r,\alpha_{r+1}+\alpha_{r+2}+2n_{r+1}+1))\\
&\quad\times\int_{-\infty}^{\infty}e^{-i\xi_{r+1}x_{r+1}}(1+\tanh x_{r+1})^{a_{r+1}}(1-\tanh x_{r+1})^{a_{r+2}}\\
&\quad\times P_{n_{r+1}}^{(\alpha_{r+2},\alpha_{r+1})}(\tanh x_{r+1})\,\mathrm{d}x_{r+1}=2^{a_{r+1}+a_{r+2}-rn_{r+1}-1}\\
&\quad\times\mathcal{F}(g_r(x_1,\ldots,x_r;n_1,\ldots,n_r,a_1,\ldots,a_r,a_{r+1}+a_{r+2}+n_{r+1},\\
&\quad\alpha_1,\ldots,\alpha_r,\alpha_{r+1}+\alpha_{r+2}+2n_{r+1}+1))\\
&\quad\times\int_0^1 t^{a_{r+1}-\frac{i\xi_{r+1}}{2}-1}(1-t)^{a_{r+2}+\frac{i\xi_{r+1}}{2}-1}P_{n_{r+1}}^{(\alpha_{r+2},\alpha_{r+1})}(2t-1)\,\mathrm{d}t\\
&=2^{a_{r+1}+a_{r+2}-rn_{r+1}-1}\mathcal{F}(g_r(x_1,\ldots,x_r;n_1,\ldots,n_r,a_1,\ldots,a_r,
\end{aligned}$$

$$
\begin{aligned}
&a_{r+1}+a_{r+2}+n_{r+1},\alpha_1,\ldots,\alpha_r,\alpha_{r+1}+\alpha_{r+2}+2n_{r+1}+1))\\
&\times\int_0^1 t^{a_{r+1}-\frac{i\xi_{r+1}}{2}-1}(1-t)^{a_{r+2}+\frac{i\xi_{r+1}}{2}-1}\frac{(\alpha_{r+2}+1)_{n_{r+1}}}{n_{r+1}!}\\
&\times {}_2F_1\left(\begin{matrix}-n_{r+1},n_{r+1}+\alpha_{r+1}+\alpha_{r+2}+1\\ \alpha_{r+2}+1\end{matrix}\,\middle|\,1-t\right)\mathrm{d}t\\
&=\frac{2^{a_{r+1}+a_{r+2}-rn_{r+1}-1}(\alpha_{r+2}+1)_{n_{r+1}}}{n_{r+1}!}\mathcal{F}\left(g_r\left(x_1,\ldots,x_r;n_1,\ldots,n_r,\right.\right.\\
&a_1,\ldots,a_r,a_{r+1}+a_{r+2}+n_{r+1},\alpha_1,\ldots,\alpha_r,\alpha_{r+1}+\alpha_{r+2}+2n_{r+1}+1)\big)\\
&\times\sum_{l=0}^{n_{r+1}}\frac{(-n_{r+1})_l(n_{r+1}+\alpha_{r+1}+\alpha_{r+2}+1)_l}{(\alpha_{r+2}+1)_l\,l!}\\
&\times\int_0^1 t^{a_{r+1}-\frac{i\xi_{r+1}}{2}-1}(1-t)^{a_{r+2}+\frac{i\xi_{r+1}}{2}-1+l}\,\mathrm{d}t\\
&=\frac{2^{a_{r+1}+a_{r+2}-rn_{r+1}-1}(\alpha_{r+2}+1)_{n_{r+1}}}{n_{r+1}!}B\left(a_{r+1}-\frac{i\xi_{r+1}}{2},\ a_{r+2}+\frac{i\xi_{r+1}}{2}\right)\\
&\times {}_3F_2\left(\begin{matrix}-n_{r+1},\ n_{r+1}+\alpha_{r+1}+\alpha_{r+2}+1,\ a_{r+2}+\frac{i\xi_{r+1}}{2}\\ \alpha_{r+2}+1,\ a_{r+1}+a_{r+2}\end{matrix}\,\middle|\,1\right)\\
&\times\mathcal{F}\left(g_r\left(x_1,\ldots,x_r;n_1,\ldots,n_r,a_1,\ldots,\right.\right.\\
&\left.\left.a_r,a_{r+1}+a_{r+2}+n_{r+1},\alpha_1,\ldots,\alpha_r,\alpha_{r+1}+\alpha_{r+2}+2n_{r+1}+1)\right),
\end{aligned}
$$

which proves the theorem.

4.2 Proof of Theorem 3.2

For the case $r=1$, the Fourier transform of the function

$$g_1\left(x_1;n_1,a_1,a_2,\alpha_1,\alpha_2\right)=(1+\tanh x_1)^{a_1}(1-\tanh x_1)^{a_2}P_{n_1}^{(\alpha_1,\alpha_2)}(\Upsilon_1),$$

is

$$
\begin{aligned}
&\mathcal{F}\left(g_1\left(x_1;n_1,a_1,a_2,\alpha_1,\alpha_2\right)\right)\\
&\quad=\int_{-\infty}^{\infty}e^{-i\xi x_1}(1+\tanh x_1)^{a_1}(1-\tanh x_1)^{a_2}P_{n_1}^{(\alpha_2,\alpha_1)}(\tanh x_1)\,\mathrm{d}x_1\\
&\quad=\frac{2^{a_1+a_2-1}(\alpha_2+1)_{n_1}}{n_1!}\Lambda_1^1(\mathbf{a},\boldsymbol{\alpha},n_1;\xi),
\end{aligned}
\tag{4.1}
$$

where

$$\Lambda_1^1(\mathbf{a},\boldsymbol{\alpha},n_1;\xi) = {}_3F_2\left(\begin{matrix}-n_1,\ a_2+\frac{i\xi}{2},\ n_1+\alpha_1+\alpha_2+1\\ \alpha_2+1,\ a_1+a_2\end{matrix}\Big|\,1\right) B\left(a_2+\frac{i\xi}{2},\ a_1-\frac{i\xi}{2}\right),$$

which was proved by Koelink [2] and it was also rewritten in terms of the continuous Hahn polynomials (2.5) as

$$\mathcal{F}\left(g_1\left(x_1;n_1,a_1,a_2,\alpha_1,\alpha_2\right)\right) = \frac{2^{a_1+a_2-1}}{i^{n_1}\left(a_1+a_2\right)_{n_1}} B\left(a_2+\frac{i\xi}{2},\ a_1-\frac{i\xi}{2}\right) \times p_{n_1}\left(\frac{\xi}{2};\ a_2,\ \alpha_1-a_1+1,\ \alpha_2-a_2+1,\ a_1\right),$$

where $\mathbf{a}=(a_1,a_2)$, $\boldsymbol{\alpha}=(\alpha_1,\alpha_2)$ and ${}_3F_2$ is a special case of the hypergeometric function (2.3). Now, let us consider the following specific function in terms of the orthogonal polynomials on 2-simplex:

$$g_2\left(x_1,x_2;n_1,n_2,a_1,a_2,a_3,\alpha_1,\alpha_2,\alpha_3\right) = \left(1+\tanh x_1\right)^{a_1}\left(1-\tanh x_1\right)^{a_2+a_3} \times \left(1+\tanh x_2\right)^{a_2}\left(1-\tanh x_2\right)^{a_3} P_{n_1,n_2}^{(\alpha_1,\alpha_2,\alpha_3)}\left(\Upsilon_1,\Upsilon_2\right), \tag{4.2}$$

where $a_1,a_2,a_3,\alpha_1,\alpha_2,\alpha_3$ are real parameters, and

$$\Upsilon_1(x_1)=\Upsilon_1=\frac{1+\tanh x_1}{2},\qquad \Upsilon_2(x_1,x_2)=\Upsilon_2=\frac{\left(1-\tanh x_1\right)\left(1+\tanh x_2\right)}{4}.$$

It is easily seen that the function g_2 can also be expressed in terms of g_1 from (3.3)

$$\begin{aligned} &g_2\left(x_1,x_2;n_1,n_2,a_1,a_2,a_3,\alpha_1,\alpha_2,\alpha_3\right)\\ &\quad= 2^{-n_2}\left(1+\tanh x_2\right)^{a_2}\left(1-\tanh x_2\right)^{a_3} P_{n_2}^{(\alpha_3,\alpha_2)}\left(\tanh x_2\right)\\ &\qquad\times g_1\left(x_1;n_1,a_1,a_2+a_3+n_2,\alpha_1,\alpha_2+\alpha_3+2n_2+1\right). \end{aligned} \tag{4.3}$$

The corresponding Fourier transform is calculated, under the substitution $\tanh x_2 = 2v-1$, as follows from:

$$\begin{aligned} &\mathcal{F}\left(g_2\left(x_1,x_2;n_1,n_2,a_1,a_2,a_3,\alpha_1,\alpha_2,\alpha_3\right)\right)\\ &= \int_{-\infty}^{\infty}\int_{-\infty}^{\infty} e^{-i\left(\xi_1x_1+\xi_2x_2\right)} g_2\left(x_1,x_2;n_1,n_2,a_1,a_2,a_3,\alpha_1,\alpha_2,\alpha_3\right)\mathrm{d}x_1\mathrm{d}x_2\\ &= 2^{-n_2}\int_{-\infty}^{\infty} e^{-i\xi_1x_1} g_1\left(x_1;n_1,a_1,a_2+a_3+n_2,\alpha_1,\alpha_2+\alpha_3+2n_2+1\right)\mathrm{d}x_1 \end{aligned}$$

$$\times \int_{-\infty}^{\infty} e^{-i\xi_2 x_2}\,(1+\tanh x_2)^{a_2}\,(1-\tanh x_2)^{a_3}\,P_{n_2}^{(\alpha_3,\alpha_2)}\,(\tanh x_2)\,\mathrm{d}x_2$$

$$= \mathcal{F}\,(g_1\,(x_1; n_1, a_1, a_2+a_3+n_2, \alpha_1, \alpha_2+\alpha_3+2n_2+1))$$

$$\times 2^{a_2+a_3-n_2-1} \int_0^1 v^{a_2-\frac{i\xi_2}{2}-1}\,(1-v)^{a_3+\frac{i\xi_2}{2}-1}\,P_{n_2}^{(\alpha_3,\alpha_2)}\,(2v-1)\,dv$$

$$= \mathcal{F}\,(g_1\,(x_1; n_1, a_1, a_2+a_3+n_2, \alpha_1, \alpha_2+\alpha_3+2n_2+1))$$

$$\times 2^{a_2+a_3-n_2-1} \int_0^1 v^{a_2-\frac{i\xi_2}{2}-1}\,(1-v)^{a_3+\frac{i\xi_2}{2}-1}\,\frac{(\alpha_3+1)_{n_2}}{n_2!}$$

$$\times\ {}_2F_1\left(\begin{matrix}-n_2, n_2+\alpha_2+\alpha_3+1\\ \alpha_3+1\end{matrix}\,\middle|\,1-v\right)dv$$

$$= \mathcal{F}\,(g_1\,(x_1; n_1, a_1, a_2+a_3+n_2, \alpha_1, \alpha_2+\alpha_3+2n_2+1))\,\frac{(\alpha_3+1)_{n_2}}{n_2!}\,2^{a_2+a_3-n_2-1}$$

$$\times \sum_{l_2=0}^{n_2} \frac{(-n_2)_{l_2}\,(n_2+\alpha_2+\alpha_3+1)_{l_2}}{(\alpha_3+1)_{l_2}\,l_2!} \int_0^1 v^{a_2-\frac{i\xi_2}{2}-1}\,(1-v)^{a_3+\frac{i\xi_2}{2}-1+l_2}\,dv,$$

that is

$$\mathcal{F}\,(g_2\,(x_1, x_2; n_1, n_2, a_1, a_2, a_3, \alpha_1, \alpha_2, \alpha_3))$$
$$= \frac{2^{a_1+2(a_2+a_3)-2}\,(2n_2+\alpha_2+\alpha_3+2)_{n_1}\,(\alpha_3+1)_{n_2}}{n_1!n_2!}\,\Lambda_1^2(\mathbf{a}, \boldsymbol{\alpha}, \mathbf{n}; \xi_1)\Lambda_2^2(\mathbf{a}, \boldsymbol{\alpha}, \mathbf{n}; \xi_2), \tag{4.4}$$

where

$$\Lambda_1^2(\mathbf{a}, \boldsymbol{\alpha}, \mathbf{n}; \xi_1) = B\left(a_1 - \frac{i\xi_1}{2}, n_2+a_2+a_3+\frac{i\xi_1}{2}\right)$$
$$\times\ {}_3F_2\left(\begin{matrix}-n_1, n_1+2n_2+\alpha_1+\alpha_2+\alpha_3+2,\ n_2+a_2+a_3+\frac{i\xi_1}{2}\\ 2n_2+\alpha_2+\alpha_3+2,\ n_2+a_1+a_2+a_3\end{matrix}\,\middle|\,1\right),$$

$$\Lambda_2^2(\mathbf{a}, \boldsymbol{\alpha}, \mathbf{n}; \xi_2) = B\left(a_2 - \frac{i\xi_2}{2}, a_3+\frac{i\xi_2}{2}\right)\ {}_3F_2\left(\begin{matrix}-n_2, n_2+\alpha_2+\alpha_3+1, a_3+\frac{i\xi_2}{2}\\ \alpha_3+1,\ a_2+a_3\end{matrix}\,\middle|\,1\right),$$

or, in terms of the continuous Hahn polynomials (2.5)

$$\Lambda_1^2(\mathbf{a}, \boldsymbol{\alpha}, \mathbf{n}; \xi_1) = \frac{n_1!B\left(a_1-\frac{i\xi_1}{2}, n_2+a_2+a_3+\frac{i\xi_1}{2}\right)}{i^{n_1}\,(n_2+a_1+a_2+a_3)_{n_1}\,(2n_2+\alpha_2+\alpha_3+2)_{n_1}}$$
$$\times\, p_{n_1}\left(\frac{\xi_1}{2}; n_2+a_2+a_3, \alpha_1-a_1+1, n_2+\alpha_2+\alpha_3-a_2-a_3+2, a_1\right),$$

$$\Lambda_2^2(\mathbf{a},\boldsymbol{\alpha},\mathbf{n};\xi_2)=\frac{n_2!B\left(a_2-\frac{i\xi_2}{2},a_3+\frac{i\xi_2}{2}\right)}{i^{n_2}\,(a_2+a_3)_{n_2}\,(\alpha_3+1)_{n_2}}\times p_{n_2}\left(\frac{\xi_2}{2};a_3,\alpha_2-a_2+1,\alpha_3-a_3+1,a_2\right),$$

where $\mathbf{a}=(a_1,a_2,a_3)$, $\boldsymbol{\alpha}=(\alpha_1,\alpha_2,\alpha_3)$, $\mathbf{n}=(n_1,n_2)$.

Using induction method, when we apply Theorem (3.1) successively, it is satisfied for each r.

4.3 Proof of Theorem 3.3

For $r=1$, by use of (3.2) and (4.1) in Parseval's identity, Koelink [2] proved that the special function

$$\begin{aligned}{}_1S_{n_1}(x_1;a_1,a_2,b_1,b_2)&={}_3F_2\left(\begin{matrix}-n_1,a_2+x_1/2,\ n_1+a_1+a_2+b_1+b_2-1\\ a_2+b_2,a_1+a_2\end{matrix}\Big|\,1\right)\\&=\frac{n_1!i^{-n_1}}{(a_1+a_2)_{n_1}\,(a_2+b_2)_{n_1}}p_{n_1}\left(\frac{-ix_1}{2};a_2,b_1,b_2,a_1\right)\end{aligned}\tag{4.5}$$

has an orthogonality relation of the form

$$\begin{aligned}&\int_{-\infty}^{\infty}\Gamma\left(a_1-\frac{ix_1}{2}\right)\Gamma\left(a_2+\frac{ix_1}{2}\right)\Gamma\left(b_1+\frac{ix_1}{2}\right)\Gamma\left(b_2-\frac{ix_1}{2}\right)\\&\quad\times\ {}_1S_{n_1}(ix_1;a_1,a_2,b_1,b_2)\ {}_1S_{m_1}(-ix_1;b_1,b_2,a_1,a_2)\,\mathrm{d}x_1\\&=\frac{2^2\pi n_1!\Gamma(n_1+a_1+b_1)\,\Gamma^2(a_2+b_2)\,\Gamma(a_1+a_2)}{(2n_1+a_1+a_2+b_1+b_2-1)\,\Gamma(n_1+a_2+b_2)}\frac{\Gamma(b_1+b_2)}{\Gamma(n_1+a_1+a_2+b_1+b_2-1)}\delta_{n_1,m_1}\\&=\frac{2^2\pi\,(n_1!)^2\ \Gamma(a_1+a_2)\,\Gamma(b_1+b_2)}{\left((a_2+b_2)_{n_1}\right)^2}h_{n_1}^{(a_1+b_1-1,a_2+b_2-1)}\delta_{n_1,m_1},\end{aligned}$$

where $h_{n_1}^{(a_1+b_1-1,a_2+b_2-1)}$ is defined as (2.8), from which it follows:

$$\begin{aligned}&\int_{-\infty}^{\infty}\Gamma(a_2+ix_1)\,\Gamma(a_1-ix_1)\,\Gamma(b_1+ix_1)\,\Gamma(b_2-ix_1)\\&\quad\times p_{n_1}(x_1;\ a_2,\ b_1,\ b_2,\ a_1)\,p_{m_1}(x_1;\ a_2,\ b_1,\ b_2,\ a_1)\,\mathrm{d}x_1\\&=\frac{2\pi\,\Gamma(n_1+a_2+b_2)\,\Gamma(n_1+a_1+b_1)\,\Gamma(n_1+a_1+a_2)\,\Gamma(n_1+b_1+b_2)}{n_1!\,(2n_1+a_1+a_2+b_1+b_2-1)\,\Gamma(n_1+a_1+a_2+b_1+b_2-1)}\delta_{n_1,m_1},\end{aligned}$$

for $a_1,a_2,b_1,b_2>0$, which gives the orthogonality relation for continuous Hahn polynomials denoted by $p_{n_1}(x_1;\ a_2,\ b_1,\ b_2,\ a_1)$.

For the case $r = 2$, by substituting (4.2) and (4.7) in Parseval's identity, we get

$$\begin{aligned}
&\int_{-\infty}^{\infty}\int_{-\infty}^{\infty} g_2\,(x_1, x_2; n_1, n_2, a_1, a_2, a_3, \alpha_1, \alpha_2, \alpha_3)\\
&\quad\times g_2\,(x_1, x_2; m_1, m_2, b_1, b_2, b_3, \beta_1, \beta_2, \beta_3)\,\mathrm{d}x_1\mathrm{d}x_2\\
&= \int_{-\infty}^{\infty}\int_{-\infty}^{\infty} (1+\tanh x_1)^{a_1+b_1}\,(1-\tanh x_1)^{a_2+a_3+b_2+b_3}\,(1+\tanh x_2)^{a_2+b_2}\\
&\quad\times (1-\tanh x_2)^{a_3+b_3}\,P_{n_1,n_2}^{(\alpha_1,\alpha_2,\alpha_3)}\,(\Upsilon_1(x_1),\,\Upsilon_2(x_1,x_2))\\
&\quad\times P_{m_1,m_2}^{(\beta_1,\beta_2,\beta_3)}\,(\Upsilon_1(x_1),\,\Upsilon_2(x_1,x_2))\,\mathrm{d}x_1\mathrm{d}x_2\\
&= \frac{1}{(2\pi)^2}\int_{-\infty}^{\infty}\int_{-\infty}^{\infty} \frac{(2n_2+\alpha_2+\alpha_3+2)_{n_1}\,(2m_2+\beta_2+\beta_3+2)_{m_1}\,(\alpha_3+1)_{n_2}}{2^{4-a_1-b_1-2(a_2+a_3+b_2+b_3)}n_1!n_2!m_1!m_2!}\\
&\quad\times(\beta_3+1)_{m_2}\,\Lambda_1^2(\mathbf{a},\boldsymbol{\alpha},\mathbf{n};\xi_1)\overline{\Lambda_1^2(\mathbf{b},\boldsymbol{\beta},\mathbf{m};\xi_1)}\Lambda_2^2(\mathbf{a},\boldsymbol{\alpha},\mathbf{n};\xi_2)\overline{\Lambda_2^2(\mathbf{b},\boldsymbol{\beta},\mathbf{m};\xi_2)}\mathrm{d}\xi_1\mathrm{d}\xi_2,
\end{aligned}$$

where $\mathbf{n} = (n_1, n_2)$, $\mathbf{m} = (m_1, m_2)$, $\boldsymbol{\alpha} = (\alpha_1, \alpha_2, \alpha_3)$, $\boldsymbol{\beta} = (\beta_1, \beta_2, \beta_3)$, $\mathbf{a} = (a_1, a_2, a_3)$ and $\mathbf{b} = (b_1, b_2, b_3)$.

Now, using the transforms $\tanh x_1 = 2u - 1$ and $\tanh x_2 = \frac{2v}{1-u} - 1$ in the left-hand side of the equality, respectively, we have

$$\begin{aligned}
&2^4\pi^2\int_0^1\int_0^{1-u} u^{a_1+b_1-1}v^{a_2+b_2-1}\,(1-u-v)^{a_3+b_3-1}\\
&\quad P_{n_1,n_2}^{(\alpha_1,\alpha_2,\alpha_3)}\,(u,v)\,P_{m_1,m_2}^{(\beta_1,\beta_2,\beta_3)}\,(u,v)\,dvdu\\
&= \int_{-\infty}^{\infty}\int_{-\infty}^{\infty} \frac{(2n_2+\alpha_2+\alpha_3+2)_{n_1}\,(2m_2+\beta_2+\beta_3+2)_{m_1}\,(\alpha_3+1)_{n_2}\,(\beta_3+1)_{m_2}}{n_1!n_2!m_1!m_2!}\\
&\quad\times\Lambda_1^2(\mathbf{a},\boldsymbol{\alpha},\mathbf{n};\xi_1)\overline{\Lambda_1^2(\mathbf{b},\boldsymbol{\beta},\mathbf{m};\xi_1)}\Lambda_2^2(\mathbf{a},\boldsymbol{\alpha},\mathbf{n};\xi_2)\overline{\Lambda_2^2(\mathbf{b},\boldsymbol{\beta},\mathbf{m};\xi_2)}\mathrm{d}\xi_1\mathrm{d}\xi_2. \quad (4.6)
\end{aligned}$$

On the other hand, if in the left-hand side of (4.6), we take $a_1 + b_1 - 1 = \alpha_1 = \beta_1$, $a_2+b_2-1 = \alpha_2 = \beta_2$, and $a_3+b_3-1 = \alpha_3 = \beta_3$, then according to the orthogonality relation (2.7), Eq. (4.6) reduces to

$$\begin{aligned}
&2^4\pi^2\int_0^1\int_0^{1-u} u^{a_1+b_1-1}v^{a_2+b_2-1}\,(1-u-v)^{a_3+b_3-1}\\
&\quad\times P_{n_1,n_2}^{(a_1+b_1-1,a_2+b_2-1,a_3+b_3-1)}\,(u,v)\,P_{m_1,m_2}^{(a_1+b_1-1,a_2+b_2-1,a_3+b_3-1)}\,(u,v)\,dvdu\\
&= 2^4\pi^2 h_{n_1,n_2}^{(a_1+b_1-1,a_2+b_2-1,a_3+b_3-1)}\delta_{n_1,m_1}\delta_{n_2,m_2}
\end{aligned}$$

$$= \int_{-\infty}^{\infty}\int_{-\infty}^{\infty} \frac{(2n_2+\alpha_2+\alpha_3+2)_{n_1}\,(2m_2+\beta_2+\beta_3+2)_{m_1}\,(\alpha_3+1)_{n_2}\,(\beta_3+1)_{m_2}}{n_1!n_2!m_1!m_2!}$$
$$\times \Lambda_1^2(\mathbf{a},\boldsymbol{\alpha},\mathbf{n};\xi_1)\overline{\Lambda_1^2(\mathbf{b},\boldsymbol{\beta},\mathbf{m};\xi_1)}\Lambda_2^2(\mathbf{a},\boldsymbol{\alpha},\mathbf{n};\xi_2)\overline{\Lambda_2^2(\mathbf{b},\boldsymbol{\beta},\mathbf{m};\xi_2)}\mathrm{d}\xi_1\mathrm{d}\xi_2,$$

so

$$\frac{2^4\pi^2\,(n_1!)^2\,(n_2!)^2\,\Gamma(n_2+a_1+a_2+a_3)\,\Gamma(n_2+b_1+b_2+b_3)\,\Gamma(a_2+a_3)\,\Gamma(b_2+b_3)}{(2n_2+a_2+b_2+a_3+b_3)_{n_1}^2\,(a_3+b_3)_{n_2}^2}$$
$$\times h_{n_1,n_2}^{(a_1+b_1-1,a_2+b_2-1,a_3+b_3-1)}\delta_{n_1,m_1}\delta_{n_2,m_2}$$
$$= \int_{-\infty}^{\infty}\int_{-\infty}^{\infty} \Theta_2(\xi_1,\xi_2;a_1,a_2,a_3)\,\Lambda_2(\xi_1,\xi_2;n_1,n_2,a_1,a_2,a_3,b_1,b_2,b_3)$$
$$\times \overline{\Theta_2(\xi_1,\xi_2;b_1,b_2,b_3)\,\Lambda_2(\xi_1,\xi_2;m_1,m_2,b_1,b_2,b_3,a_1,a_2,a_3)}\mathrm{d}\xi_1\mathrm{d}\xi_2,$$

where

$$\Theta_2(\xi_1,\xi_2;a_1,a_2,a_3)$$
$$= \Gamma\left(a_1-\frac{i\xi_1}{2}\right)\Gamma\left(a_2+a_3+\frac{i\xi_1}{2}\right)\Gamma\left(a_2-\frac{i\xi_2}{2}\right)\Gamma\left(a_3+\frac{i\xi_2}{2}\right)\left(a_2+a_3+\frac{i\xi_1}{2}\right)_{n_2},$$
$$\Lambda_2(\xi_1,\xi_2;n_1,n_2,a_1,a_2,a_3,b_1,b_2,b_3)$$
$$= {}_3F_2\left(\begin{matrix}-n_2,\ n_2+a_2+b_2+a_3+b_3-1,\ a_3+\frac{i\xi_2}{2}\\ a_3+b_3,\ a_2+a_3\end{matrix}\,\middle|\,1\right)$$
$$\times {}_3F_2\left(\begin{matrix}-n_1,\ n_1+2n_2+a_1+b_1+a_2+b_2+a_3+b_3-1,\ n_2+a_2+a_3+\frac{i\xi_1}{2}\\ 2n_2+a_2+b_2+a_3+b_3,\ n_2+a_1+a_2+a_3\end{matrix}\,\middle|\,1\right),$$

or, in terms of the Hahn polynomials (2.5)

$$\Lambda_2(\xi_1,\xi_2;n_1,n_2,a_1,a_2,a_3,b_1,b_2,b_3)$$
$$= \frac{n_1!n_2!i^{-n_1-n_2}}{(a_2+a_3)_{n_2}\,(a_3+b_3)_{n_2}\,(n_2+a_1+a_2+a_3)_{n_1}\,(2n_2+a_2+b_2+a_3+b_3)_{n_1}}$$
$$\times p_{n_1}\left(\frac{\xi_1}{2};n_2+a_2+a_3,b_1,n_2+b_2+b_3,a_1\right)p_{n_2}\left(\frac{\xi_2}{2};a_3,b_2,b_3,a_2\right),$$

and $h_{n_1,n_2}^{(\alpha_1,\alpha_2,\alpha_3)}$ is given by (2.8). By applying the method in the cases $r=1$ and $r=2$, if we substitute (3.1) and (3.7) in the Parseval's identity (3.8), after the necessary calculations, we obtain the desired result.

4.4 Proof of Proposition 3.6

We first consider the case $r=1$. Substituting $m_1 \to -n_1$, $m_2 \to n_1+a_1+a_2+b_1+b_2-1$, $m_3 \to a_2+\frac{x_1}{2}$, $s_1 \to a_2+b_2$, $s_2 \to a_1+a_2$ and $z \to 1$ in the relation

(3.10) and taking into account the relation

$$ {}_1S_{n_1}(x_1;a_1,a_2,b_1,b_2) = {}_3F_2\left(\begin{matrix}-n_1,\ a_2+x_1/2,\ n_1+a_1+a_2+b_1+b_2-1\\ a_2+b_2,\ a_1+a_2\end{matrix}\Big|\,1\right) $$

given by (4.5), we obtain the recurrence relation for the family of the special functions ${}_1S_{n_1}(x_1;a_1,a_2,b_1,b_2)$

$$ \begin{aligned}&(B_1+C_1)\ {}_1S_{n_1}(x_1;a_1,a_2,b_1,b_2)+(B_2+C_2)\ {}_1S_{n_1+1}(x_1;a_1,a_2,b_1-1,b_2)\\ &\quad+B_3\ {}_1S_{n_1+2}(x_1;a_1,a_2,b_1-2,b_2)=0,\end{aligned} \tag{4.7} $$

where

$$ \begin{aligned}B_1+C_1&=\frac{b_1-1+\frac{x_1}{2}}{n_1},\quad B_3=-\frac{(n_1+a_2+b_2+1)\,(n_1+a_1+a_2+1)}{(n_1)_2},\\ B_2+C_2&=\frac{(a_2+b_2)\,(a_1+a_2)-(n_1+a_1+a_2+b_1+b_2-1)\left(a_2+\frac{x_1}{2}\right)}{(n_1)_2}\\ &\quad+\frac{n_1+a_1+2a_2-b_1+b_2+2-\frac{x_1}{2}}{n_1}.\end{aligned} $$

For $r=2$, the relation (3.13) gives the result

$$ \begin{aligned}&{}_2S_{n_1,n_2}(x_1,x_2;a_1,a_2,a_3,b_1,b_2,b_3)\\ &\quad={}_3F_2\left(\begin{matrix}-n_2,\ n_2+a_2+a_3+b_2+b_3-1,\ a_3+\frac{x_2}{2}\\ a_3+b_3,\ a_2+a_3\end{matrix}\Big|\,1\right)\\ &\qquad\times\left(a_2+a_3+\frac{x_1}{2}\right)_{n_2}\ {}_1S_{n_1}(x_1;a_1,a_2+a_3+n_2,b_1,b_2+b_3+n_2)\,.\end{aligned} \tag{4.8} $$

If the equation obtained by taking $b_2\to n_2+b_2+b_3$ ve $a_2\to n_2+a_2+a_3$ in relation (4.7) is multiplied by

$$ \left(a_2+a_3+\frac{x_1}{2}\right)_{n_2}\ {}_3F_2\left(\begin{matrix}-n_2,\ n_2+a_2+a_3+b_2+b_3-1,\ a_3+\frac{x_2}{2}\\ a_3+b_3,\ a_2+a_3\end{matrix}\Big|\,1\right) $$

and the relation (4.8) is used, we get

$$ \begin{aligned}&\frac{b_1-1+\frac{x_1}{2}}{n_1}\ {}_2S_{n_1,n_2}(x_1,x_2;a_1,a_2,a_3,b_1,b_2,b_3)\\ &\quad+\left(\frac{(n_2+a_1+a_2+a_3)\,(2n_2+a_2+a_3+b_2+b_3)}{(n_1)_2}\right.\\ &\quad\left.-\frac{(n_1+2n_2+a_1+a_2+a_3+b_1+b_2+b_3-1)\left(n_2+a_2+a_3+\frac{x_1}{2}\right)}{(n_1)_2}\right.\end{aligned} $$

$$+\frac{n_1+3n_2+a_1+2\,(a_2+a_3)-b_1+b_2+b_3+2-\frac{x_1}{2}}{n_1}\Bigg)$$
$$\times\ {}_2S_{n_1+1,n_2}\,(x_1,x_2;a_1,a_2,a_3,b_1-1,b_2,b_3)$$
$$-\frac{(n_1+2n_2+a_2+a_3+b_2+b_3+1)\,(n_1+n_2+a_1+a_2+a_3+1)}{(n_1)_2}$$
$$\times\ {}_2S_{n_1+2,n_2}\,(x_1,x_2;a_1,a_2,a_3,b_1-2,b_2,b_3)=0.$$

Similarly, if relation (4.7) is applied consecutively in view of the relation (3.13), then the statement of Proposition (3.6) is obtained.

4.5 Proof of Proposition 3.7

By taking $m_1\to n_1+a_1+a_2+b_1+b_2-1, m_2\to -n_1, m_3\to a_2+\frac{x_1}{2}, s_1\to a_2+b_2$, $s_2\to a_1+a_2$ and $z\to 1$ in the relation (3.10), we obtain

$$(B_1+C_1)\ {}_1S_{n_1}\,(x_1;a_1,a_2,b_1,b_2)+(B_2+C_2)\ {}_1S_{n_1}\,(x_1;a_1,a_2,b_1-1,b_2)$$
$$+B_3\ {}_1S_{n_1}\,(x_1;a_1,a_2,b_1-2,b_2)=0, \tag{4.9}$$

where

$$B_1+C_1=-\frac{b_1-1+\frac{x_1}{2}}{n_1+a_1+a_2+b_1+b_2-1},$$
$$B_3=-\frac{(n_1+a_1+b_1-2)\,(n_1+b_1+b_2-2)}{(n_1+a_1+a_2+b_1+b_2-2)_2},$$
$$B_2+C_2=\frac{(a_2+b_2)\,(a_1+a_2)+n_1\left(a_2+\frac{x_1}{2}\right)}{(n_1+a_1+a_2+b_1+b_2-2)_2}+\frac{n_1-a_2+2b_1-3+\frac{x_1}{2}}{n_1+a_1+a_2+b_1+b_2-1}$$

for the case $r=1$. If we apply the recurrence relation (4.9) successively, we obtain the required result in view of the relation (3.14).

4.6 Proof of Proposition 3.8

For the case $r=1$, if we get $m_1\to a_2+\frac{x_1}{2}, m_2\to -n_1, m_3\to n_1+a_1+a_2+b_1+b_2-1, s_1\to a_2+b_2, s_2\to a_1+a_2$ and $z\to 1$ in relation (3.10), the following relation holds for the family of the special function ${}_1S_{n_1}\,(x_1;a_1,a_2,b_1,b_2)$

$$(B_1+C_1)\ {}_1S_{n_1}\,(x_1;a_1,a_2,b_1,b_2)-(B_2+C_2)\ {}_1S_{n_1}\,(x_1;a_1+1,a_2-1,$$
$$b_1-1,b_2+1)+B_3\ {}_1S_{n_1}\,(x_1;a_1+2,a_2-2,b_1-2,b_2+2)=0,$$

where

$$B_1+C_1=\left(b_1-1+\frac{x_1}{2}\right)\left(a_2-1+\frac{x_1}{2}\right),$$
$$B_3=\left(a_1+1-\frac{x_1}{2}\right)\left(b_2+1-\frac{x_1}{2}\right),$$

$$B_2 + C_2 = (a_2 + b_2)(a_1 + a_2) + n_1(n_1 + a_1 + a_2 + b_1 + b_2 - 1) - (a_1 + a_2 - b_1 + b_2 + 2 - x_1)\left(a_2 - 1 + \frac{x_1}{2}\right).$$

It is enough to consider the relation (3.14) and use the last relation successively to obtain the statement of Proposition (3.8).

4.7 Proof of Proposition 3.9

Taking $m_1 \to -n_1, m_2 \to n_1 + a_1 + a_2 + b_1 + b_2 - 1, m_3 \to a_2 + \frac{x_1}{2}, s_1 \to a_1 + a_2$, $s_2 \to a_2 + b_2$ and $z \to 1$ in (3.11), we arrive at the following relation for the case $r = 1$:

$$(B_1 + C_1)\ {}_1S_{n_1}(x_1; a_1, a_2, b_1, b_2) + (B_2 + C_2)\ {}_1S_{n_1}(x_1; a_1 + 1, a_2, b_1 - 1, b_2) + C_3\ {}_1S_{n_1}(x_1; a_1 + 2, a_2, b_1 - 2, b_2) = 0,$$

where

$$B_1 + C_1 = -\frac{b_1 - 1 + \frac{x_1}{2}}{a_1 + a_2 - 1},$$

$$C_3 = -\frac{(n_1 + a_1 + a_2 + 1)(n_1 + b_1 + b_2 - 2)\left(a_1 + 1 - \frac{x_1}{2}\right)}{(a_1 + a_2 - 1)_3},$$

$$B_2 + C_2 = \frac{(2(a_1 + a_2) + 1)\left(a_2 + b_1 + b_2 - 2 + \frac{x_1}{2}\right)}{(a_1 + a_2 - 1)_2} + \frac{n_1(n_1 + a_1 + a_2 + b_1 + b_2 - 1)}{(a_1 + a_2 - 1)_2} - \frac{(a_1 + a_2 + b_1 + b_2 - 1)\left(a_2 + \frac{x_1}{2}\right)}{(a_1 + a_2 - 1)_2} - \frac{a_2 + b_2 - 1}{a_1 + a_2 - 1}.$$

If this relation is applied consecutively in (3.13), the proof is completed.

4.8 Proof of Proposition 3.10

For the case $r = 1$, if we replace $m_1 \to -n_1, m_2 \to n_1 + a_1 + a_2 + b_1 + b_2 - 1$, $m_3 \to a_2 + \frac{x_1}{2}, s_1 \to a_2 + b_2, s_2 \to a_1 + a_2$ and $z \to 1$ in relation (3.11), the family of the special function ${}_1S_{n_1}(x_1; a_1, a_2, b_1, b_2)$ satisfies the relation

$$(B_1 + C_1)\ {}_1S_{n_1}(x_1; a_1, a_2, b_1, b_2) + (B_2 + C_2)\ {}_1S_{n_1}(x_1; a_1, a_2, b_1 - 1, b_2 + 1) + C_3\ {}_1S_{n_1}(x_1; a_1, a_2, b_1 - 2, b_2 + 2) = 0,$$

where

$$B_1 + C_1 = -\frac{b_1 - 1 + \frac{x_1}{2}}{a_2 + b_2 - 1},$$

$$C_3 = -\frac{(n_1+a_2+b_2+1)\,(n_1+a_1+b_1-2)\left(b_2+1-\frac{x_1}{2}\right)}{(a_2+b_2-1)_3},$$

$$\begin{aligned} B_2+C_2 = {} & \frac{(2\,(a_2+b_2)+1)\left(a_1+a_2+b_1-2+\frac{x_1}{2}\right)}{(a_2+b_2-1)_2} \\ & + \frac{n_1\,(n_1+a_1+a_2+b_1+b_2-1)}{(a_2+b_2-1)_2} \\ & - \frac{(a_1+a_2+b_1+b_2-1)\left(a_2+\frac{x_1}{2}\right)}{(a_2+b_2-1)_2} - \frac{a_1+a_2-1}{a_2+b_2-1}. \end{aligned}$$

Applying this relation consecutively in (3.14) provides to obtain the statement of Proposition (3.10).

4.9 Proof of Proposition 3.11

By taking $m_1 \to n_1+a_1+a_2+b_1+b_2-1, m_2 \to -n_1, m_3 \to a_2+\frac{x_1}{2}, s_1 \to a_1+a_2,$ $s_2 \to a_2+b_2$ and $z \to 1$ in relation (3.12), we find the recurrence relation for the family of the special function ${}_1S_{n_1}\,(x_1; a_1, a_2, b_1, b_2)$ for the case $r=1$

$$\begin{aligned} & (a_1+a_2)\ {}_1S_{n_1}\,(x_1; a_1, a_2, b_1, b_2) \\ & \quad + (n_1+b_1+b_2-1)\ {}_1S_{n_1}\,(x_1; a_1+1, a_2, b_1-1, b_2) \\ & \quad - (n_1+a_1+a_2+b_1+b_2-1)\ {}_1S_{n_1}\,(x_1; a_1+1, a_2, b_1, b_2) = 0. \end{aligned}$$

If we use the last relation consecutively by taking into account (3.14), it is seen that the family of the special function ${}_rS_{\mathbf{n}}\,(\mathbf{x}; \mathbf{a}, \mathbf{b})$ satisfies the desired recurrence relation for $r \geq 1$.

Acknowledgements The authors would like to thank the three reviewers for their many valuable suggestions and comments which led us to improve this paper.

Author Contributions All authors have contributed equally to this work. All authors have read and approved the final manuscript.

Funding Open Access funding provided thanks to the Universidade de Vigo/CISUG agreement with Springer Nature. The work of R.A. has been partially supported by TUBITAK Research Grant Proj. No. 120F140. The work of I.A. has been partially supported by the Agencia Estatal de Investigación (AEI) of Spain under Grant PID2020-113275GB-I00, cofinanced by the European Community fund FEDER.

Data availability Not applicable.

Declarations

Conflict of interest The authors declare that they have no competing interests.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included

in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

References

1. Erdélyi, A., Magnus, W., Oberhettinger, F., et al.: Tables of Integral Transforms, vol. II. McGraw-Hill Book Company, Inc., New York (1954)
2. Koelink, H.T.: On Jacobi and continuous Hahn polynomials. Proc. Am. Math. Soc. **124**(3), 887–898 (1996)
3. Koepf, W., Masjed-Jamei, M.: Two classes of special functions using Fourier transforms of some finite classes of classical orthogonal polynomials. Proc. Am. Math. Soc. **135**(11), 3599–3606 (2007)
4. Koornwinder, T.H.: Group Theoretic Interpretations of Askey's Scheme of Hypergeometric Orthogonal Polynomials. In: Orthogonal polynomials and their applications (Segovia, 1986). (Lecture Notes in Math.; Vol. 1329), pp. 46–72. Springer, Berlin (1988)
5. Koornwinder, T.H.: Special Orthogonal Polynomial Systems Mapped Onto Each Other by the Fourier-Jacobi Transform. In: Orthogonal polynomials and applications (Bar-le-Duc, 1984). (Lecture Notes in Math.; Vol. 1171), pp. 174–183, Springer, Berlin (1985)
6. Masjed-Jamei, M., Koepf, W.: Two classes of special functions using Fourier transforms of generalized ultraspherical and generalized Hermite polynomials. Proc. Am. Math. Soc. **140**(6), 2053–2063 (2012)
7. Masjed-Jamei, M., Marcellán, F., Huertas, E.J.: A finite class of orthogonal functions generated by Routh–Romanovski polynomials. Complex Var. Elliptic Equ. **59**(2), 162–171 (2014)
8. Masjed-Jamei, M., Koepf, W.: Two finite classes of orthogonal functions. Appl. Anal. **92**(11), 2392–2403 (2013)
9. Koepf, W., Masjed-Jamei, M.: A generic polynomial solution for the differential equation of hypergeometric type and six sequences of orthogonal polynomials related to it. Integral Transf. Spec. Funct. **17**(8), 559–576 (2006)
10. Masjed-Jamei, M.: Three finite classes of hypergeometric orthogonal polynomials and their application in functions approximation. Integral Transf. Spec. Funct. **13**(2), 169–191 (2002)
11. Masjed-Jamei, M.: A basic class of symmetric orthogonal polynomials using the extended Sturm–Liouville theorem for symmetric functions. J. Math. Anal. Appl. **325**(2), 753–775 (2007)
12. Tratnik, M.V.: Some multivariable orthogonal polynomials of the Askey tableau-continuous families. J. Math. Phys. **32**(8), 2065–2073 (1991)
13. Tratnik, M.V.: Some multivariable orthogonal polynomials of the Askey tableau-discrete families. J. Math. Phys. **32**(9), 2337–2342 (1991)
14. Koelink, H.T., Van der Jeugt, J.: Convolutions for orthogonal polynomials from Lie and quantum algebra representations. SIAM J. Math. Anal. **29**, 794–822 (1998)
15. Güldoğan, E., Aktaş, R., Area, I.: Some classes of special functions using Fourier transforms of some two-variable orthogonal polynomials. Integral Transf. Spec. Funct. **31**(6), 437–470 (2020)
16. Koornwinder, T.H.: Two-variable analogues of the classical orthogonal polynomials. In: Askey, R.A. (ed.) Theory and Application of Special Functions. Proceedings of an Advanced Seminar, pp. 435–495. Academic Press, New York (1975)
17. Fernández, L., Pérez, T.E., Piñar, M.A.: On Koornwinder classical orthogonal polynomials in two variables. J. Comput. Appl. Math. **236**(15), 3817–3826 (2012)
18. Rainville, E.D.: Special Functions. Macmillan, New York (1960)
19. Szegő, G.: Orthogonal Polynomials. Amer. Math. Soc. Colloq. Publ., vol. 23, 4th ed. Am. Math. Soc., Providence (1975)
20. Abramowitz, M., Stegun, I.A. (editors): Handbook of mathematical functions with formulas, graphs, and mathematical tables. Wiley., New York (1984). A Wiley-Interscience Publication; reprint of the 1972 edition
21. Askey, R.: Continuous Hahn polynomials. J. Phys. A **18**(16), L1017–L1019 (1985)
22. Alhaidari, A.D., Li, Y.T.: Quantum Systems Associated with the Hahn and Continuous Hahn Polynomials. Rep. Math. Phys. **82**(3), 285–301 (2018)

23. Dunkl, C.F., Xu, Y.: Orthogonal polynomials of several variables. 2nd ed. (Encyclopedia of Mathematics and its Applications; Vol. 155). Cambridge University Press, Cambridge (2014)
24. Davies, B.: Integral transforms and their applications. 3rd ed. (Texts in Applied Mathematics; Vol. 41). Springer, New York (2002)

25. Petkovšek, M., Wilf, H.S., Zeilberger, D.: $A = B$. A K Peters, Ltd., Wellesley, Mass (1996)
26. Fasenmyer, C.: A Note on Pure Recurrence Relations. Am. Math. Monthly. **56**(1), 14–17 (1949)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.