

Lack of evidence for improved β -adrenoceptor-mediated papillary muscle contraction by low-dose empagliflozin treatment in a rat model of streptozotocin-induced diabetes

Zeynep Elif YESILYURT-DIRICAN^{1,2} , Irem KARAOMERLIOGLU^{1,5} , Betül Rabia ERDOGAN-VADACCA¹ , Gaye OZTURK^{1,3} , Martin Christian MICHEL⁴ , Ebru ARIOGLU-INAN^{1,*} 

¹ Department of Pharmacology, Faculty of Pharmacy, Ankara University, 06560, Ankara, Türkiye

² Department of Pharmacology, Faculty of Pharmacy, Gazi University, 06330, Ankara, Türkiye

³ Department of Pharmacology, Faculty of Pharmacy, Hacettepe University, 06100, Ankara, Türkiye

⁴ Department of Pharmacology, University Medical Center, Johannes Gutenberg University, 55131, Mainz, Germany

⁵ present affiliation; Turkish Medicines and Medical Devices Agency

* Corresponding Author. E-mail: arioglu@ankara.edu.tr (E.A.I.); Tel. +90-312-203 31 45.

Received: 19 November 2024 / Revised: 20 March 2025 / Accepted: 23 March 2025

ABSTRACT: Diabetes mellitus leads to cardiovascular complications including impaired cardiac β -adrenoceptor (β -AR) function. Sodium-glucose cotransporter-2 (SGLT2) inhibitors, such as empagliflozin (EMPA), improve outcomes in heart failure patients and animal models. Therefore, we have investigated the effects of EMPA on in vivo cardiac function and β -AR mediated contractile responses in streptozotocin (STZ)-induced diabetes in a design reflecting late-onset of treatment. Male Sprague Dawley rats were divided into 4 groups (control, EMPA-treated control, diabetic, and EMPA-treated diabetic). Diabetes was induced by STZ injection (40 mg/kg). 13-16 weeks after STZ injection, a low dose of EMPA (10 mg/kg/day, daily oral gavage) or vehicle was administered for another 8 weeks. At the end of the treatment period, in vivo cardiac function was evaluated by pressure-volume (PV) loop analysis and β -AR mediated contractile response was determined by isoprenaline on isolated papillary strips. The blood glucose-lowering effect of low-dose EMPA was confirmed. EMPA did not change cardiac function in control rats. Diabetic rats had a reduced heart rate, cardiac output, stroke work, rate of contraction and rate of relaxation and increased isovolumic relaxation, whereas in vitro responses were not markedly attenuated. Treatment with EMPA showed a trend for improvement of some (e.g., stroke volume, ejection fraction, cardiac index) but not all parameters. Our results indicate that low-dose EMPA treatment had limited effects on cardiac impairment despite reducing blood glucose when initiated after diabetes has manifested. Future studies using a preventive rather than therapeutic approach could help to clarify the possible benefits of EMPA on the diabetic heart.

KEYWORDS: β -adrenoceptor; diabetes; empagliflozin; heart ; inotropy; pressure-volume loop analysis.

1. INTRODUCTION

Diabetes remains an important health problem despite the steps taken to better understand its pathophysiology and new treatment approaches [1]. The International Diabetes Federation estimates that it affected 451 million people worldwide in 2017 and will affect 693 million by 2045 [2]. Cardiovascular complications are one of the most important causes of morbidity and mortality in patients with type 1 and type 2 diabetes (T1DM and T2DM, respectively) [3]. While good glycemic control is important to prevent microvascular diabetic complications such as nephropathy, retinopathy and neuropathy, the ACCORD [4], ADVANCE [5] and VADT [6] trials have shown that major cardiovascular events, considered to represent macrovascular complications, were inadequately addressed by treatment approaches despite a tight glycemic control.

Inhibitors of the sodium-glucose cotransporter-2 (SGLT2) are antidiabetic drugs that inhibit glucose reabsorption in the renal tubules to stimulate glucose excretion [7]. The cardiovascular outcome studies of empagliflozin (EMPA) [8], canagliflozin [9], dapagliflozin [10] and ertugliflozin [11] have shown that

How to cite this article: Yeşilyurt-Dirican ZE, Karaomerlioglu I, Erdogan-Vadacca BR, Ozturk G, Michel MC, Arioglu-Inan E. Lack of evidence for improved β -adrenoceptor-mediated papillary muscle contraction by low-dose empagliflozin treatment in a rat model of streptozotocin-induced diabetes. J Res Pharm. 2026; 30(1): 60-71.

diabetic patients treated with SGLT2 inhibitors had a lower risk of major cardiovascular events compared with placebo including a reduced rate of hospitalization for heart failure and lower all-cause mortality. These studies sparked additional trials designed to study effects on heart failure that included diabetic and non-diabetic patients, either in heart failure with reduced [12, 13] or preserved ejection fraction [14, 15]. SGLT2 inhibitors effectively reduced heart failure endpoints in such studies, and that effect was similarly pronounced in heart failure patients with and without concomitant diabetes. Accordingly, SGLT2 inhibitors are now recommended by major guidelines for the routine management of heart failure patients [16,17].

The effects of SGLT2 inhibitors have been explored in various rodent models of diabetes-associated cardiomyopathy. These have used T2DM models including ob/ob mice [18, 19], db/db mice with or without additional angiotensin II infusion [20, 21], rats [22-24] and mice [25] treated with a combination of a high-fat diet and injection of streptozotocin (STZ), Zucker diabetic fatty rats [26], seipin knock-out mice [27]), a cross-breed of mRen27 and Tet29 rats [28], and Dahl salt-sensitive rats on a high-sodium diet [29]. They have used various SGLT2 inhibitors including canagliflozin, dapagliflozin and EMPA. While SGLT2 inhibitors are only indicated for the treatment of T2DM, some studies have also tested them in animal models of T1DM, i.e., rats and, less frequently, mice injected with STZ [30-35]. Moreover, canagliflozin had been shown to improve oxidative stress in the heart in an euglycemic model in which cardiomyopathy had been induced by infusion of the β -AR agonist isoprenaline [36]. These studies generally confirmed the clinically observed beneficial effect on diabetic cardiomyopathy; however, most of them applied a preventive or early intervention design (starting treatment soon after onset of disease), which is unlikely to occur in patients [37].

Despite the compelling evidence for the use of SGLT2 inhibitors in the treatment of heart failure, the underlying mechanisms remain to be understood. Given that SGLT2 inhibitors are similarly effective in the treatment of heart failure in patients with and without diabetes [12-15], their beneficial effects on heart failure in non-diabetic patients indicate that they are not a direct consequence of their glucose lowering effects. While the clinical studies yielded limited mechanistic explanation of the beneficial effects, the above rodent studies have proposed a range of potential molecular and cellular mechanisms that could contribute including those related to the inflammasome, fibroblast activation or activation of various protein kinases.

The probably most consistently observed cellular and molecular phenotype in heart failure, in general, is a decreased responsiveness to β -adrenoceptor (β -AR) stimulation that is at least partly due to down-regulation of the receptor [38-41]. Both desensitization and down-regulation primarily affect β_1 -AR, whereas β_2 -AR are less consistently affected, and β_3 -AR even are up-regulated in most studies [42]. The regulation of cardiac β -AR in animal models of T1DM has been explored in more than 50 studies; while most have reported a down-regulation of β_1 - and β_2 -AR, this was not always observed and even an increased expression was reported in some cases; whether down-regulation was observed was not obviously linked to duration of diabetes for periods up to about 30 weeks [43]. Animal studies show that desensitization and down-regulation of β -AR, particularly β_1 -AR, also occurs in the diabetic heart, e.g., in rats injected with STZ [44, 45] or in Zucker diabetic fatty rats [46]. Surprisingly, to the best of our knowledge, the effect of SGLT2 inhibitors on reduced β -AR responses in (diabetic) cardiomyopathy has not been investigated.

Therefore, the present pilot study was designed to explore effects of EMPA as example for SGLT2 inhibitors in general on β -AR responses as assessed in isolated papillary muscle in a rat model of STZ-induced diabetes in a therapeutic setting, i.e., when treatment is started several weeks after the induction of diabetes; effects on general cardiac function were assessed by pressure-volume (PV) loop analysis. For this purpose, we have applied a dose of EMPA that is in the low range of reportedly effective doses in rats [34, 47, 48]

2. RESULTS

2.1. General animal characteristics

STZ injection markedly increased blood glucose, which was substantially attenuated but not fully abolished by EMPA (Figure 1), thereby validating the induction of diabetes by STZ and the antidiabetic efficacy of low-dose EMPA in this model. While the control and EMPA-treated control groups had similar blood glucose and body weight, diabetic rats had a reduced body weight that was not affected by EMPA (Figure 1). Heart weight was similar across all groups leading to a slightly increased heart/body weight ratio in diabetic and EMPA-treated diabetic rats (Figure 1).




Figure 1. General characteristics of the rats at study end. Blood glucose, body weight, heart weight and heart/body weight ratio in control (C; n=7), EMPA-treated control (CE; n=6), diabetic (D; n=7) and EMPA-treated diabetic rats (DE; n=9). Each data point represents one animal, bars represent means $\pm$ SD. A statistical analysis was not performed because these measurements are intended for model validation only.

2.2. In vivo baseline hemodynamic parameters

Systolic, diastolic, and mean blood pressure were measured when the PV loop catheter was inserted in the carotid artery. The values were slightly reduced in the diabetic groups, but no improvement was observed in EMPA treated diabetic group (Table 1). Cardiac function parameters were assessed by using the PV loop catheter and most of these parameters are summarized in Table 1.

Table 1. In vivo basal hemodynamic parameters in control (C; n=7), EMPA-treated control (CE; n=5), diabetic (D; n=6) and EMPA-treated diabetic rats (DE; n=8).

	C	CE	D	DE
SBP, mm Hg	133 $\pm$ 32	123 $\pm$ 11	114 $\pm$ 16	116 $\pm$ 13
DBP, mm Hg	95 $\pm$ 25	96 $\pm$ 16	85 $\pm$ 7	90 $\pm$ 10
MAP, mm Hg	120 $\pm$ 29	114 $\pm$ 12	105 $\pm$ 13	107 $\pm$ 12
HR, beats/min	285 $\pm$ 25	282 $\pm$ 11	250 $\pm$ 22	242 $\pm$ 19
CO, ml/min	55 $\pm$ 12	58 $\pm$ 9	39 $\pm$ 5	49 $\pm$ 8
SW, mm Hg/ μ l	22 $\pm$ 4	19 $\pm$ 3	16 $\pm$ 3	19 $\pm$ 4
EDV, μ l	388 $\pm$ 28	374 $\pm$ 50	357 $\pm$ 70	391 $\pm$ 53
ESV, μ l	193 $\pm$ 24	167 $\pm$ 26	199 $\pm$ 45	188 $\pm$ 29
SV, μ l	195 $\pm$ 38	207 $\pm$ 30	157 $\pm$ 30	203 $\pm$ 34
EDVI, μ l/g 10^3	0.91 $\pm$ 0.09	0.90 $\pm$ 0.15	1.03 $\pm$ 0.20	1.14 $\pm$ 0.19
ESVI, μ l/g 10^3	0.45 $\pm$ 0.08	0.40 $\pm$ 0.57	0.57 $\pm$ 0.12	0.55 $\pm$ 0.10
CI, ml/min g 10^3	129 $\pm$ 27	141 $\pm$ 23	113 $\pm$ 15	143 $\pm$ 26
SVI, μ l/g 10^3	0.45 $\pm$ 0.09	0.50 $\pm$ 0.08	0.46 $\pm$ 0.10	0.59 $\pm$ 0.12

Values are presented as means $\pm$ SD after administering isoflurane anesthesia. SBP: Systolic blood pressure; DBP: diastolic blood pressure; MAP: mean arterial pressure HR: heart rate; CO: Cardiac output; SW: Stroke work; ESV: End-systolic volume; EDV: End-diastolic volume; SV: Stroke volume; EDVI: End-diastolic volume index; ESVI: End-systolic volume index; CI: cardiac index; SVI: stroke volume index. A statistical analysis was not performed because these measurements are intended for model validation only.

The parameters closely related to systolic and diastolic function are shown in Figure 2 and 3. Our results indicate that some of the hemodynamic parameters such as heart rate, cardiac output and stroke work were impaired in the diabetic group; however, EMPA treatment caused only minor if any improvement of these parameters in diabetics. Other parameters were found to be similar in all four groups (Table 1). The systolic function parameters end-systolic pressure, rate of contraction and ejection fraction decreased in the diabetic group. These parameters were slightly greater in EMPA-treated diabetics

compared to diabetic group (Figure 2). On the other hand, EMPA had little effect on systolic function in control animals.




Figure 2. In vivo basal systolic parameters. End-systolic pressure, rate of contraction (+dp/dt) and ejection fraction in control (C; n=7), EMPA-treated control (CE; n=5), diabetic (D; n=6) and EMPA-treated diabetic rats (DE; n=8). Each data point represents one animal, bars represent means $\pm$ SD. A statistical analysis was not performed because these measurements are intended for model validation only.

Similarly, diastolic function was impaired in the diabetic group. EMPA showed a slight improvement, however, was unable to normalize the relaxation rate, end-diastolic pressure or the time constant of isovolumic relaxation (Tau) in diabetics (Figure 3).




Figure 3. In vivo basal diastolic parameters. End-diastolic pressure, rate of relaxation (-dp/dt) and isovolumic relaxation time constant (Tau) in control (C; n=7), EMPA-treated control (CE; n=5), diabetic (D; n=6) and EMPA-treated diabetic rats (DE; n=8). Each data point represents one animal, bars represent means $\pm$ SD. A statistical analysis was not performed because these measurements are intended for model validation only.

Preload independent parameters (preload recruitable stroke work, end-systolic pressure-volume relationship, end-diastolic pressure-volume relationship) were measured by vena cava inferior occlusions for 5 seconds (Table 2). The expected deterioration was not observed in the diabetic group, making it futile to comment on a possible effect of EMPA in diabetic animals.

Table 2. Preload independent in vivo cardiac parameters end-systolic pressure-volume relationship (ESPVR), end-diastolic pressure-volume relationship (EDPVR) and preload recruitable stroke work (PRSW) in control (C; n=7), EMPA-treated control (CE; n=5), diabetic (D; n=5) and EMPA-treated diabetic rats (DE; n=7). Values are presented as means $\pm$ SD. A statistical analysis was not performed because these measurements are intended for model validation only.

	C	CE	D	DE
ESPVR	0.29 $\pm$ 0.08	0.33 $\pm$ 0.15	0.28 $\pm$ 0.06	0.32 $\pm$ 0.10
EDPVR	0.005 $\pm$ 0.004	0.006 $\pm$ 0.003	0.008 $\pm$ 0.005	0.008 $\pm$ 0.006
PRSW	53 $\pm$ 8	58 $\pm$ 11	56 $\pm$ 11	53 $\pm$ 6

2.3. Contractile responses to isoprenaline in papillary muscle

The responses to β -AR agonist isoprenaline (0.1 nM-10 μ M) were evaluated in the papillary muscle isolated from the left ventricle. Although the inotropic response in diabetics seems to have decreased slightly compared to controls (Figure 4), this decrease was lower than expected. The contraction response in the EMPA group was like that in the control group. On the other hand, treatment with EMPA did not ameliorate the decreased contractility due to diabetes (Figure 4). The p-value for the overall ANOVA ($p=0.0913$, $F=2.490$, $df=3$) was insufficient to reject the null hypothesis of inter-group differences. The corresponding -log

EC₅₀ values in the four groups were also similar across groups (C, 7.26±0.29; CE 7.23±0.19; D: 7.22±0.19; DE: 7.32±0.31).




Figure 4. Responses to isoprenaline in papillary muscle. Concentration-response curves and maximal responses to isoprenaline in control (C; n=4), EMPA-treated control (CE; n=5), diabetic (D; n=6) and EMPA-treated diabetic rats (CE; n=8). For curves only mean values are shown for clarity, whereas each data point represents one animal and bars represent means ± SD for calculated E_{max}. The p-value for the overall ANOVA was 0.0913.

3. DISCUSSION

SGLT2 inhibitors have shown beneficial effects in heart failure patients with and without concomitant diabetes [12-15]. Desensitization of cardiac β -AR is probably the most consistent pathophysiological finding at the cellular and molecular level in heart failure [38-41], and also occurs in diabetic animals [44-46]; however, effects of SGLT2 inhibitors on cardiac β -AR function have not been reported to the best of our knowledge. Therefore, a pilot study was performed to explore the effects of the SGLT2 inhibitor EMPA on responses to the β -AR agonist isoprenaline in the rat STZ model of diabetes.

3.1. Critique of methods

SGLT2 inhibitors are approved for the treatment of T2DM and at least some (dapagliflozin and EMPA) irrespective of diabetic status also for the treatment of heart failure. Their effects on the heart have been studied in various rodent models of T2DM [18-23, 25-29]. Although SGLT2 inhibitors are not used for the treatment of T1DM, some studies have explored their cardiac effects in STZ-based rat and mouse models thereof [30-33], where they also exhibited beneficial effects. The rationale for this is based on the mechanism of action of SGLT2 inhibitors, i.e., promoting renal glucose excretion that can be useful in both T2DM and T1DM patients. Accordingly, some clinical studies have demonstrated beneficial effects of SGLT2 inhibitors in T1DM patients [49-51]. We have used an STZ-based model T1DM. The key advantage of this is that it typically produces a more severe diabetic and cardiovascular phenotype than most models of T2DM, which facilitates the detection of therapeutic effects. Clinical studies with heart failure patients with reduced [12, 13] or preserved ejection fraction [14, 15] have shown that SGLT2 inhibitors are similarly beneficial in those with and without concomitant diabetes. Therefore, it should not matter for an experimental study exploring the effect of SGLT2 inhibitors on cardiomyopathy whether diabetes was present, whether it was T1DM or T2DM or whether it was induced by STZ or other means. Against this background, we have used the STZ-induced diabetes model because it tends to yield a more severe cardiac phenotype than T2DM models.

Previous studies in the rat STZ model have largely been based on doses of 30-60 mg/kg for rats (considerably higher doses were used in most mouse studies) [52], and studies in diabetic cardiomyopathy typically have also used similar doses [30, 31, 34]. Our laboratory has previously used STZ doses of 32-45 mg/kg [53-59]; these exhibited a phenotype with cardiac impairment including diminished β -AR mediated contractility of isolated papillary muscle [56]. We prefer these lower doses because they tend to cause less mortality than e.g., 50-60 mg/kg [60]. Therefore, the present study was based on an STZ dose of 40 mg/kg, which caused a major increase in blood glucose. Despite our hope to reduce attrition due to mortality by choosing a low STZ dose, we have observed a greater loss of animals than in our previous studies [53-59], leading to smaller sample sizes than intended.

Our experiments have initiated EMPA treatment 13-16 weeks after STZ injection, i.e., at a time point when diabetic cardiomyopathy can be expected to have fully developed. This is later than most other studies in the field, which started SGLT2 inhibitor treatment in most cases ≤ 1 week after STZ injection [22, 24, 25, 30, 32-35], and only rarely at later time points such as 2 weeks [31] or 8 weeks after STZ injection [23]. We

consider our approach to have higher translational validity as the clinical diagnosis of T2DM typically is made 2-7 years after the onset of the disease [37]. On the other hand, structural cardiac damage including fibrosis [23, 31] may have fully developed at such late time points, which could make it more difficult for any medication to reverse this. In other words, most previous studies in this field have used study designs that are largely preventive based on the early start of the intervention, whereas ours and one other study [23] belong to the very few applying a therapeutic design, i.e., starting the SGLT2 inhibitor treatment after cardiomyopathy has fully developed. Of note, SGLT2 inhibitors have proven beneficial in clinical studies [8-11], i.e., under conditions of initiation of treatment considerably after onset of diabetes [37].

Meta-analyses have shown that the beneficial effects of dapagliflozin and EMPA in heart failure patients are quantitatively similar [61] and major international guidelines assume this to be a class effect [16, 17]. Rodent studies with EMPA have mostly used daily oral doses of 10-30 mg/kg [7], and studies related to diabetic cardiomyopathy have also used those doses [21, 28, 30, 31, 34]. While both doses appeared effective, the only direct comparative study in the field indicates that the daily 10 mg/kg dose used by us produces a less pronounced blood glucose lowering and improvement of cardiac function [34]. Our study used crushed EMPA tablets for the treatments, and the reduced blood glucose levels confirmed that this resulted in an efficacious delivery of the treatment. While we cannot exclude that the excipients in the clinically used material have contributed to that, we consider this highly unlikely because these excipients are generally considered to be inert. In a previous study crushed dapagliflozin tablets had also effectively decreased blood glucose [62]. As discussed by those authors, using tablets may enhance the translational value of the study as it better mimics the clinical condition. Various approaches have been used to assess cardiac function in rodent models of diabetes. While the most frequently used technique is echocardiography [21-23, 25-29, 32] others including PV loop analysis [19, 30] cardiac catheters in general [34], ECG analysis [27], heart weight [20, 24, 33] and cardiac enzymes such as LDH and CK-MB have also been used [35]. Our *in vivo* studies are based on PV loop analysis, which like other approaches using left ventricular catheterization is a robust method to determine systolic and diastolic function.

3.2. Model validation

Our blood glucose data demonstrate the expected elevation upon treatment with STZ and lowering with EMPA (Figure 1). This was associated with reductions of end-systolic pressure and cardiac contractility and relaxation *in vivo* (Figure 2 and 3) and reduced contractile responses in the papillary muscle preparations (Figure 4). Accordingly, our study in principle yielded the expected phenotype of diabetic cardiomyopathy. However, the degree of impairment was smaller than what is normally seen by us [53-59] and others in the rat STZ model [31-35]. This smaller than expected severity of cardiomyopathy can potentially be at least partly explained by the unexpectedly high mortality and may reflect that the surviving rats had a less severe cardiac phenotype than the overall group.

3.3. Effects of EMPA on diabetic cardiomyopathy

Our experiments found a trend for improvement of cardiac function upon administration of EMPA to diabetic rats that was consistent ejection fraction, relaxation, cardiac output, stroke work, stroke volume or cardiac index but not for many other parameters including responses to the β -AR agonist isoprenaline. A reduced expression of cardiac β_1 - and β_2 -AR has been observed often in animal models of T1DM, but lack of major alterations has also been reported repeatedly [43], the latter observation being in line with the lack of major changes of isoprenaline responses in the present study. While the latter findings are not conclusive based on small sample size and large variability, they make it unlikely that SGLT2 inhibitor treatment has a major effect on the desensitization of cardiac β -AR in heart failure.

The overall moderate effects of EMPA in the present study were rather disappointing. Part of this could possibly be explained by lower sample sizes than planned due to an unexpectedly high attrition and to a less severe cardiac phenotype than in our previous studies using similar STZ doses [53-59], which in turn may at least partly reflect that animals with a more severe phenotype may preferentially have contributed to the attrition. Both factors would markedly reduce the statistical power to detect the beneficial effects of EMPA.

However, there also is a possible biological reason for the weaker effects of EMPA in the present as compared to previous studies using this or other SGLT2 inhibitors in models involving STZ injections with or without a prior/concomitant high-fat diet [22, 24, 25, 30-35]). As noted above, these studies started administration of the SGLT2 inhibitor mostly ≤ 1 week after the STZ injection, i.e., at a time point when structural damage to the heart may yet have been rudimentary or even absent. The present data do not allow

to address this possibility. However, it is noteworthy that other studies from our group using STZ injection found that EMPA prevents a diabetic phenotype in the urinary bladder when administration starts early (prevention design) [63] but not when it starts late (treatment design) [64]. Given that T2DM is often diagnosed several years after onset, our findings may differ from those of others but nonetheless more reflective of the clinical situation. This is a testable hypothesis for future studies.

4. CONCLUSION

In summary, the present pilot study does not provide robust evidence for a beneficial effect of EMPA in the STZ injection-based rat model of T1DM when administration of EMPA started 13-16 weeks after STZ injection, i.e., in a therapeutic setting. This contradicts other studies reporting beneficial effects of SGLT2 inhibitors including EMPA in STZ-based diabetes model with or without a prior/concomitant high-fat diet. However, those studies initiated the treatment with an SGLT2 inhibitor much earlier (mostly within ≤ 1 week after STZ injection), which may reflect a preventive and not a therapeutic setting. While we cannot exclude that other factors including a surprisingly small severity of diabetic cardiomyopathy and surprisingly large attrition due to mortality have contributed to our negative data, we propose that future studies should explicitly compare early/preventive and late/therapeutic settings when evaluating the effects of SGLT2 inhibitors in diabetic cardiomyopathy. In this sense, our pilot study provided what a pilot study should do, it led to a novel testable hypothesis. Moreover, our data may assist in the assumptions required to generate robust sample size/power calculations for such studies.

5. MATERIALS AND METHODS

5.1. Animals and treatments

Male Sprague Dawley rats (11-12 weeks old) were obtained from Bilkent University, Department of Molecular Biology and Genetics Animal Unit (Ankara, Türkiye) and housed under 12-h light/dark cycle (7 a.m. to 7 p.m.) at $22 \pm 1^\circ\text{C}$ ambient temperature. All experimental procedures of the present study were in line with NIH guidelines for the care and use of experimental animals and had been approved by the Animal Welfare Committee of Ankara University (06/02/2019; 2019-4-41). The rats had free access to standard rat chow (Purina Rat Chow; Optima AS, Bolu, Türkiye) and tap water. After one week of acclimatization, the rats were randomly divided into control and diabetic groups. We set the sample sizes as 14 rats for the control group and 20 rats for the diabetic groups to account for possible attrition in the diabetic groups because of a greater mortality risk. Diabetes was induced by intraperitoneal (*ip*) injection of STZ (40 mg/kg dissolved in citrate buffer at pH 4.5). Control rats were injected with citrate buffer. Three days after STZ injection, blood glucose level was measured and rats with blood glucose levels higher than 300 mg/dl were considered as diabetic. If blood glucose levels were lower than 300 mg/dl, a second or third STZ injection was administered as commonly done in this model; in the present study, this affected two rats each in the diabetic and in diabetic+EMPA group.

The treatment started at 13-16 weeks after STZ injection. For this purpose, EMPA (Jardiance®) tablets were crushed and suspended in distilled water. Each tablet includes lactose monohydrate, microcrystalline cellulose, hydroxypropyl cellulose, croscarmellose sodium, colloidal anhydrous silica, magnesium stearate. The film coating of the tablets also contains hypromellose, titanium dioxide, talc, macrogol, and yellow ferric oxide. The crushed material was dissolved in 5 ml of distilled water. Control and diabetic rats were randomly divided into 2 groups each to yield control, EMPA-treated control, diabetic and EMPA-treated diabetic groups. One rat in the diabetic group was euthanized due to poor health condition in this period. Thus, the sample size became 7 in each control group, 8 in diabetic and 11 in EMPA-treated diabetic group. The sample size was higher in the treated diabetic group regarding the possible mortality risk. Treated control and diabetic rats received 10 mg/kg EMPA (calculated for active pharmaceutical ingredient) by once daily oral gavage for 8 weeks. The dose was selected based on previous studies, where the lower but effective of two doses under investigation was 10 mg/kg/day [34, 47, 48]. The control and diabetic rats received distilled water by oral gavage. In the treatment period, one rat from the EMPA-treated control group was killed during the gavage procedure. One rat in the diabetic group returned to normoglycemic level and was excluded from the study. Two rats in the EMPA-treated diabetic group were euthanized due to tumor development in week 7. At the end of the treatment period, the rats were sacrificed, and *in vivo* and *in vitro* experiments were performed (control $n=7$, EMPA-treated control $n=6$, diabetic $n=7$ and EMPA-treated diabetic $n=9$).

5.2. In vivo measurement of basal hemodynamic parameters

PV loop analysis was performed as previously reported [56]. The rats were anesthetized by 2% isoflurane inhalation and body temperature was monitored to be 37°C with a rectal probe during the operation. After incision, a PV catheter (Transonic, London, Ontario, Canada) was inserted into the right carotid artery. After 3-minute of blood pressure recording, the catheter was advanced into the left ventricle. Following a 10-minute stabilization period, PV loops were recorded. Then, preload-independent cardiac parameters were measured by 5-second vena cava occlusions. At the end of the experiments, the following basal hemodynamic parameters were calculated; end-diastolic pressure, end-systolic pressure, heart rate, end-diastolic volume, end-systolic volume, cardiac output, ejection fraction, stroke volume, stroke work, rate of contraction, rate of relaxation, isovolumic relaxation constant (Tau logistic) and preload independent parameters; preload recruitable stroke work, end-systolic pressure-volume relationship, end-diastolic pressure-volume relationship. End-systolic volume, end-diastolic volume, stroke volume and cardiac output were normalized to body weight to eliminate the differences of body weight between animals (end-systolic volume index, end-diastolic volume index, stroke volume index, cardiac index) [65].

5.3. In vitro papillary muscle experiments

The experiments were performed as previously described [56]. Briefly, the rats were anesthetized under 2% isoflurane inhalation. After PV loop analysis, hearts were rapidly isolated. The extracted hearts were transferred to an oxygenated experimental plate containing heparin and Krebs solution (120 mM NaCl; 4.8 mM KCl; 1.25 mM $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$; 1.25 mM $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$; 1.2 mM KH_2PO_4 ; 25 mM NaHCO_3 and 10 mM glucose at pH 7.4). Papillary muscles were dissected from the left ventricle and mounted in horizontal organ baths (Harvard Apparatus GmbH, March-Hugstetten, Germany). Electrical field stimulation (0.6 Hz, 2 ms, twice the diastolic threshold) was applied to the muscle [66] and the organ bath was perfused with Krebs solution (95% O_2 ; 5% CO_2 ; 30°C) at a rate of 5 ml/min. Tension was recorded using a force transducer (HSE F30; Type 372, Harvard Apparatus GmbH). Following a 60-minute stabilization period, the maximal tension ($L_{\max}$) was found by stretching the papillary muscles with 10 μm increments and the experiments were then performed at 90% of $L_{\max}$. After equilibration, cumulative concentration-response curves of isoprenaline (0.1 nM-10 μM) were obtained.

5.4. Chemicals

EMPA tablets (Jardiance®, 25 mg, Boehringer Ingelheim) were purchased from a local pharmacy. The ingredients of Krebs solution, STZ and isoprenaline were obtained from Sigma Aldrich (Darmstadt, Germany). Isoflurane was from Adeka (Samsun, Türkiye). A Bayer glucometer and glucose test strips (Contour Plus, Switzerland) were used to measure blood glucose levels.

5.5. Data analysis

The efficacy of isoprenaline in the papillary muscle experiments was determined by fitting a 3-parameter model (top, bottom, log EC_{50}) of concentration-response curve to the experimental data from each experiment to derive an estimated maximum effect ($E_{\max}$). Data are expressed as mean $\pm$ SD and shown as bar graphs overlaid with scatter plots for increased transparency [67]. To avoid misleading y-axis scaling, all y-axes start at 0 [68]. All statistical analyses were performed with Prism (version 9, GraphPad, La Jolla, CA, USA). We did not perform pre-study sample size/power calculations because meaningful sample size/power calculations require robust evidence-based assumptions on minimum biologically relevant effect sizes and on expected variability. However, we considered the pre-study evidence on the primary outcome parameter (cardiac β -AR function) to be insufficient for such estimates. Therefore, the study was designed as a pilot study to provide signals, not as a hypothesis-testing study. As the study was of exploratory nature and in line with recent recommendations [69, 70], the obtained data were not interpreted as hypothesis-testing but only as descriptive. We did not perform p-value calculations for the parameters intended for model validation but only for our target parameter, the $E_{\max}$ for isoprenaline in the isolated papillary muscle experiments. For this purpose, ANOVA assuming a normal distribution was performed; as this did not yield a low p-value, no post-tests were performed for intergroup comparisons.

Acknowledgements: This research was funded by grants from the Scientific and Technological Research Council of Türkiye (TUBITAK SBAG-119S769). Z.E.Y.D. was supported by TUBITAK (2211/A) during her PhD studies. G.O. was an undergraduate student supported by TUBITAK (119S769).

Author contributions: Conceptualization, E.A.I. and Z.E.Y.D.; formal analysis, M.C.M.; investigation, E.A.I., Z.E.Y.D., I.K., B.R.E. and G.O.; data curation, M.C.M.; writing-original draft preparation, E.A.I. and Z.E.Y.D.; writing-review and editing, M.C.M., E.A.I., Z.E.Y.D.; visualization, M.C.M.; supervision, E.A.I.; project administration, E.A.I.; funding

acquisition, E.A.I. All authors have read and agreed to the final version of the manuscript.

Conflict of interest statement: M.C.M. is a consultant to Boehringer Ingelheim. All other authors declare no conflict of interest. The funder had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

REFERENCES

- [1] Pennig J, Scherrer P, Gissler MC, Anto-Michel N, Hoppe N, Fünér L, Härdtnér C, Stachon P, Wolf D, Hilgendorf I, Mullick A, Bode C, Zirlik A, Goldberg IJ, Willecke F. Glucose lowering by SGLT2-inhibitor empagliflozin accelerates atherosclerosis regression in hyperglycemic STZ-diabetic mice. *Sci Rep*. 2019; 9(1):17937. <https://doi.org/10.1038/s41598-019-54224-9>
- [2] Cho NH, Shaw JE, Karuranga S, Huang Y, da Rocha Fernandes JD, Ohlrogge AW, Malanda B. IDF Diabetes Atlas: Global estimates of diabetes prevalence for 2017 and projections for 2045. *Diabetes Res Clin Pract*. 2018; 138:271-281. <https://doi.org/10.1016/j.diabres.2018.02.023>
- [3] Lee TI, Chen YC, Lin YK, Chung CC, Lu YY, Kao YH, Chen YJ. Empagliflozin Attenuates Myocardial Sodium and Calcium Dysregulation and Reverses Cardiac Remodeling in Streptozotocin-Induced Diabetic Rats. *Int J Mol Sci*. 2019 Apr 4;20(7):1680. <https://doi.org/10.3390/ijms20071680>
- [4] Action to Control Cardiovascular Risk in Diabetes Study Group; Gerstein HC, Miller ME, Byington RP, Goff DC Jr, Bigger JT, Buse JB, Cushman WC, Genuth S, Ismail-Beigi F, Grimm RH Jr, Probstfield JL, Simons-Morton DG, Friedewald WT. Effects of intensive glucose lowering in type 2 diabetes. *N Engl J Med*. 2008;358(24):2545-2559. <https://doi.org/10.1056/NEJMoa0802743>
- [5] ADVANCE Collaborative Group; Patel A, MacMahon S, Chalmers J, Neal B, Billot L, Woodward M, Marre M, Cooper M, Glasziou P, Grobbee D, Hamet P, Harrap S, Heller S, Liu L, Mancía G, Mogensen CE, Pan C, Poultier N, Rodgers A, Williams B, Bompoint S, de Galan BE, Joshi R, Travert F. Intensive blood glucose control and vascular outcomes in patients with type 2 diabetes. *N Engl J Med*. 2008;358(24):2560-2572. <http://dx.doi.org/10.1056/NEJMoa0802987>
- [6] Duckworth W, Abraira C, Moritz T, Reda D, Emanuele N, Reaven PD, Zieve FJ, Marks J, Davis SN, Hayward R, Warren SR, Goldman S, McCarren M, Vitek ME, Henderson WG, Huang GD; VADT Investigators. Glucose control and vascular complications in veterans with type 2 diabetes. *N Engl J Med*. 2009;360(2):129-139. doi: 10.1056/NEJMoa0808431. Epub 2008 Dec 17. Erratum in: *N Engl J Med*. 2009;361(10):1028. Erratum in: *N Engl J Med*. 2009;361(10):1024-1025. <https://doi.org/10.1056/NEJMoa0808431>
- [7] Michel MC, Mayoux E, Vallon V. A comprehensive review of the pharmacodynamics of the SGLT2 inhibitor empagliflozin in animals and humans. *Naunyn Schmiedeberg's Arch Pharmacol*. 2015; 388(8):801-816. <https://doi.org/10.1007/s00210-015-1134-1>
- [8] Zinman B, Wanner C, Lachin JM, Fitchett D, Bluhmki E, Hantel S, Mattheus M, Devins T, Johansen OE, Woerle HJ, Broedl UC, Inzucchi SE; EMPA-REG OUTCOME Investigators. Empagliflozin, Cardiovascular Outcomes, and Mortality in Type 2 Diabetes. *N Engl J Med*. 2015; 373(22):2117-2128. <https://doi.org/10.1056/NEJMoa1504720>
- [9] Neal B, Perkovic V, Mahaffey KW, de Zeeuw D, Fulcher G, Erondú N, Shaw W, Law G, Desai M, Matthews DR; CANVAS Program Collaborative Group. Canagliflozin and Cardiovascular and Renal Events in Type 2 Diabetes. *N Engl J Med*. 2017; 377(7):644-657. <https://doi.org/10.1056/NEJMoa1611925>
- [10] Wiviott SD, Raz I, Bonaca MP, Mosenzon O, Kato ET, Cahn A, Silverman MG, Zelniker TA, Kuder JF, Murphy SA, Bhatt DL, Leiter LA, McGuire DK, Wilding JPH, Ruff CT, Gause-Nilsson IAM, Fredriksson M, Johansson PA, Langkilde AM, Sabatine MS; DECLARE-TIMI 58 Investigators. Dapagliflozin and Cardiovascular Outcomes in Type 2 Diabetes. *N Engl J Med*. 2019; 380(4):347-357. <https://doi.org/10.1056/NEJMoa1812389>
- [11] Cannon CP, Pratley R, Dagogo-Jack S, Mancuso J, Huyck S, Masiukiewicz U, Charbonnel B, Frederich R, Gallo S, Cosentino F, Shih WJ, Gantz I, Terra SG, Cherney DZI, McGuire DK; VERTIS CV Investigators. Cardiovascular outcomes with ertugliflozin in type 2 diabetes. *N Engl J Med*. 2020; 383(15):1425-1435. <https://doi.org/10.1056/NEJMoa2004967>
- [12] McMurray JJV, Solomon SD, Inzucchi SE, Køber L, Kosiborod MN, Martinez FA, Ponikowski P, Sabatine MS, Anand IS, Bělohávek J, Böhm M, Chiang CE, Chopra VK, de Boer RA, Desai AS, Diez M, Drozd J, Dukát A, Ge J, Howlett JG, Katova T, Kitakaze M, Ljungman CEA, Merkely B, Nicolau JC, O'Meara E, Petrie MC, Vinh PN, Schou M, Tereshchenko S, Verma S, Held C, DeMets DL, Docherty KF, Jhund PS, Bengtsson O, Sjöstrand M, Langkilde AM; DAPA-HF Trial Committees and Investigators. Dapagliflozin in Patients with Heart Failure and Reduced Ejection Fraction. *N Engl J Med*. 2019;381(21):1995-2008. <https://doi.org/10.1056/NEJMoa1911303>
- [13] Packer M, Anker SD, Butler J, Filippatos G, Pocock SJ, Carson P, Januzzi J, Verma S, Tsutsui H, Brueckmann M, Jamal W, Kimura K, Schnee J, Zeller C, Cotton D, Bocchi E, Böhm M, Choi DJ, Chopra V, Chuquiere E, Giannetti N, Janssens S, Zhang J, Gonzalez Juanatey JR, Kaul S, Brunner-La Rocca HP, Merkely B, Nicholls SJ, Perrone S, Pina I, Ponikowski P, Sattar N, Senni M, Seronde MF, Spinar J, Squire I, Taddei S, Wanner C, Zannad F; EMPEROR-Reduced Trial Investigators. Cardiovascular and Renal Outcomes with Empagliflozin in Heart Failure. *N Engl J Med*. 2020;383(15):1413-1424. <https://doi.org/10.1056/NEJMoa2022190>

- [14] Anker SD, Butler J, Filippatos G, Ferreira JP, Bocchi E, Böhm M, Brunner-La Rocca HP, Choi DJ, Chopra V, Chuquiere-Valenzuela E, Giannetti N, Gomez-Mesa JE, Janssens S, Januzzi JL, Gonzalez-Juanatey JR, Merkely B, Nicholls SJ, Perrone SV, Piña IL, Ponikowski P, Senni M, Sim D, Spinar J, Squire I, Taddei S, Tsutsui H, Verma S, Vinereanu D, Zhang J, Carson P, Lam CSP, Marx N, Zeller C, Sattar N, Jamal W, Schnaidt S, Schnee JM, Brueckmann M, Pocock SJ, Zannad F, Packer M; EMPEROR-Preserved Trial Investigators. Empagliflozin in Heart Failure with a Preserved Ejection Fraction. *N Engl J Med.* 2021;385(16):1451-1461. <https://doi.org/10.1056/NEJMoa2107038>
- [15] Solomon SD, McMurray JJV, Claggett B, de Boer RA, DeMets D, Hernandez AF, Inzucchi SE, Kosiborod MN, Lam CSP, Martinez F, Shah SJ, Desai AS, Jhund PS, Belohlavek J, Chiang CE, Borleffs CJW, Comin-Colet J, Dobreanu D, Drozd J, Fang JC, Alcocer-Gamba MA, Al Habeeb W, Han Y, Cabrera Honorio JW, Janssens SP, Katova T, Kitakaze M, Merkely B, O'Meara E, Saraiva JFK, Tereshchenko SN, Thierer J, Vaduganathan M, Vardeny O, Verma S, Pham VN, Wilderäng U, Zozerska N, Bachus E, Lindholm D, Petersson M, Langkilde AM; DELIVER Trial Committees and Investigators. Dapagliflozin in Heart Failure with Mildly Reduced or Preserved Ejection Fraction. *N Engl J Med.* 2022;387(12):1089-1098. <https://doi.org/10.1056/NEJMoa2206286>
- [16] McDonagh TA, Metra M, Adamo M, Gardner RS, Baumbach A, Böhm M, Burri H, Butler J, Čelutkienė J, Chioncel O, Cleland JGF, Coats AJS, Crespo-Leiro MG, Farmakis D, Gilard M, Heymans S, Hoes AW, Jaarsma T, Jankowska EA, Lainscak M, Lam CSP, Lyon AR, McMurray JJV, Mebazaa A, Mindham R, Muneretto C, Francesco Piepoli M, Price S, Rosano GMC, Ruschitzka F, Kathrine Skibelund A; ESC Scientific Document Group. 2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure. *Eur Heart J.* 2021;42(36):3599-3726. doi: 10.1093/eurheartj/ehab368. Erratum in: *Eur Heart J.* 2021;42(48):4901. <https://doi.org/10.1093/eurheartj/ehab368>
- [17] Writing Committee; Maddox TM, Januzzi JL Jr, Allen LA, Breathett K, Butler J, Davis LL, Fonarow GC, Ibrahim NE, Lindenfeld J, Masoudi FA, Motiwala SR, Oliveros E, Patterson JH, Walsh MN, Wasserman A, Yancy CW, Youmans QR. 2021 Update to the 2017 ACC Expert Consensus Decision Pathway for Optimization of Heart Failure Treatment: Answers to 10 Pivotal Issues About Heart Failure With Reduced Ejection Fraction: A Report of the American College of Cardiology Solution Set Oversight Committee. *J Am Coll Cardiol.* 2021;77(6):772-810. <https://doi.org/10.1016/j.jacc.2020.11.022>
- [18] Ye Y, Bajaj M, Yang HC, Perez-Polo JR, Birnbaum Y. SGLT-2 inhibition with dapagliflozin reduces the activation of the Nlrp3/ASC inflammasome and attenuates the development of diabetic cardiomyopathy in mice with type 2 diabetes. Further augmentation of the effects with saxagliptin, a DPP4 inhibitor. *Cardiovasc Drugs Ther.* 2017; 31(2):119-132. <https://doi.org/10.1007/s10557-017-6725-2>
- [19] Hammoudi N, Jeong D, Singh R, Farhat A, Komajda M, Mayoux E, Hajjar R, Lebeche D. Empagliflozin Improves Left Ventricular Diastolic Dysfunction in a Genetic Model of Type 2 Diabetes. *Cardiovasc Drugs Ther.* 2017;31(3):233-246. <https://doi.org/10.1007/s10557-017-6734-1>
- [20] Arow M, Waldman M, Yadin D, Nudelman V, Shainberg A, Abraham NG, Freimark D, Kornowski R, Aravot D, Hochhauser E, Arad M. Sodium-glucose cotransporter 2 inhibitor Dapagliflozin attenuates diabetic cardiomyopathy. *Cardiovasc Diabetol.* 2020;19(1):7. <https://doi.org/10.1186/s12933-019-0980-4>
- [21] Habibi J, Aroor AR, Sowers JR, Jia G, Hayden MR, Garro M, Barron B, Mayoux E, Rector RS, Whaley-Connell A, DeMarco VG. Sodium glucose transporter 2 (SGLT2) inhibition with empagliflozin improves cardiac diastolic function in a female rodent model of diabetes. *Cardiovasc Diabetol.* 2017;16(1):9. <https://doi.org/10.1186/s12933-016-0489-z>
- [22] Liu L, Luo H, Liang Y, Tang J, Shu Y. Dapagliflozin ameliorates STZ-induced cardiac hypertrophy in type 2 diabetic rats by inhibiting the calpain-1 expression and nuclear transfer of NF-κB. *Comput Math Methods Med.* 2022; 2022:3293054. <https://doi.org/10.1155/2022/3293054>
- [23] Tian J, Zhang M, Suo M, Liu D, Wang X, Liu M, Pan J, Jin T, An F. Dapagliflozin alleviates cardiac fibrosis through suppressing EndMT and fibroblast activation via AMPKα/TGF-β/Smad signalling in type 2 diabetic rats. *J Cell Mol Med.* 2021;25(16):7642-7659. <https://doi.org/10.1111/jcmm.16601>
- [24] El-Sayed N, Mostafa YM, AboGresha NM, Ahmed AAM, Mahmoud IZ, El-Sayed NM. Dapagliflozin attenuates diabetic cardiomyopathy through erythropoietin up-regulation of AKT/JAK/MAPK pathways in streptozotocin-induced diabetic rats. *Chem Biol Interact.* 2021; 347:109617. <https://doi.org/10.1016/j.cbi.2021.109617>
- [25] Sun P, Wang Y, Ding Y, Luo J, Zhong J, Xu N, Zhang Y, Xie W. Canagliflozin attenuates lipotoxicity in cardiomyocytes and protects diabetic mouse hearts by inhibiting the mTOR/HIF-1α pathway. *iScience.* 2021;24(6):102521. <https://doi.org/10.1016/j.isci.2021.102521>
- [26] Lim VG, Bell RM, Arjun S, Kolatsi-Joannou M, Long DA, Yellon DM. SGLT2 inhibitor, canagliflozin, attenuates myocardial infarction in the diabetic and nondiabetic heart. *ACC Basic Transl Sci.* 2019; 4(1):15-26. <https://doi.org/10.1016/j.jacbs.2018.10.002>
- [27] Joubert M, Jagu B, Moutaigne D, Marechal X, Tesse A, Ayer A, Dollet L, Le May C, Toumaniantz G, Manrique A, Charpentier F, Staels B, Magré J, Cariou B, Prieur X. The Sodium-Glucose Cotransporter 2 Inhibitor Dapagliflozin Prevents Cardiomyopathy in a Diabetic Lipodystrophic Mouse Model. *Diabetes.* 2017;66(4):1030-1040. <https://doi.org/10.2337/db16-0733>

- [28] Kräker K, Herse F, Golic M, Reichhart N, Crespo-Garcia S, Strauß O, Grune J, Kintscher U, Ebrahim M, Bader M, Alenina N, Heuser A, Luft FC, Müller DN, Dechend R, Haase N. Effects of empagliflozin and target-organ damage in a novel rodent model of heart failure induced by combined hypertension and diabetes. *Sci Rep*. 2020;10(1):14061. <https://doi.org/10.1038/s41598-020-70708-5>
- [29] Ma S, He LL, Zhang GR, Zuo QJ, Wang ZL, Zhai JL, Zhang TT, Wang Y, Ma HJ, Guo YF. Canagliflozin mitigates ferroptosis and ameliorates heart failure in rats with preserved ejection fraction. *Naunyn Schmiedebergs Arch Pharmacol*. 2022;395(8):945-962. <https://doi.org/10.1007/s00210-022-02243-1>
- [30] Ideishi A, Suematsu Y, Tashiro K, Morita H, Kuwano T, Tomita S, Nakai K, Miura SI. Combination of Linagliptin and Empagliflozin Preserves Cardiac Systolic Function in an Ischemia-Reperfusion Injury Mice With Diabetes Mellitus. *Cardiol Res*. 2021;12(2):91-97. <https://doi.org/10.14740/cr1194>
- [31] Trang NN, Chung CC, Lee TW, Cheng WL, Kao YH, Huang SY, Lee TL, Chen YJ. Empagliflozin and Liraglutide Differentially Modulate Cardiac Metabolism in Diabetic Cardiomyopathy in Rats. *Int J Mol Sci*. 2021;22(3):1177. <https://doi.org/10.3390/ijms22031177>
- [32] Xing YJ, Liu BH, Wan SJ, Cheng Y, Zhou SM, Sun Y, Yao XM, Hua Q, Meng XJ, Cheng JH, Zhong M, Zhang Y, Lv K, Kong X. A SGLT2 Inhibitor Dapagliflozin Alleviates Diabetic Cardiomyopathy by Suppressing High Glucose-Induced Oxidative Stress in vivo and in vitro. *Front Pharmacol*. 2021 ;12:708177. <https://doi.org/10.3389/fphar.2021.708177>
- [33] Hodrea J, Saeed A, Molnar A, Fintha A, Barcsi A, Wagner LJ, Szabo AJ, Fekete A, Balogh DB. SGLT2 inhibitor dapagliflozin prevents atherosclerotic and cardiac complications in experimental type 1 diabetes. *PLoS One*. 2022;17(2):e0263285. <https://doi.org/10.1371/journal.pone.0263285>
- [34] Zhou Y, Wu W. The sodium-glucose co-transporter 2 inhibitor, empagliflozin, protects against diabetic cardiomyopathy by inhibition of the endoplasmic reticulum stress pathway. *Cell Physiol Biochem*. 2017; 41(6):2503-2512. <https://doi.org/10.1159/000475942>
- [35] El-Shafey M, El-Agawy MSE, Eldosoky M, Ebrahim HA, Elsherbini DMA, El-Sherbiny M, Asseri SM, Elsherbiny NM. Role of Dapagliflozin and Liraglutide on Diabetes-Induced Cardiomyopathy in Rats: Implication of Oxidative Stress, Inflammation, and Apoptosis. *Front Endocrinol (Lausanne)*. 2022;13:862394. <https://doi.org/10.3389/fendo.2022.862394>
- [36] Hasan R, Lasker S, Hasan A, Zerín F, Zamila M, Chowdhury FI, Nayan SI, Rahman MM, Khan F, Subhan N, Alam MA. Canagliflozin attenuates isoprenaline-induced cardiac oxidative stress by stimulating multiple antioxidant and anti-inflammatory signaling pathways. *Sci Rep*. 2020 ;10(1):14459. <https://doi.org/10.1038/s41598-020-71449-1>
- [37] National Institute for Health and Care Excellence. Type 2 diabetes in adults: management 2019 <https://www.nice.org.uk/guidance/ng28> (accessed December 2019).
- [38] Harding SE, Brown LA, Wynne DG, Davies CH, Poole-Wilson PA. Mechanisms of β adrenoceptor desensitisation in the failing human heart. *Cardiovasc Res*. 1994; 28:1451-1460. <https://doi.org/10.1093/cvr/28.10.1451>
- [39] Brodde OE, Michel MC. Adrenergic and muscarinic receptors in the human heart. *Pharmacol. Rev*. 1999; 51(4):651-689. [https://doi.org/10.1016/S0031-6997\(24\)01425-X](https://doi.org/10.1016/S0031-6997(24)01425-X)
- [40] Brodde OE, Bruck H, Leineweber K. Cardiac adrenoceptors: physiological and pathophysiological relevance. *J Pharmacol Sci*. 2006; 100(5):323-337. <https://doi.org/10.1254/jphs.CRJ06001X>
- [41] Lohse MJ, Engelhardt S, Eschenhagen T. What is the role of beta-adrenergic signaling in heart failure? *Circ Res*. 2003; 93(10):896-906. <https://doi.org/10.1161/01.RES.0000102042.83024.CA>
- [42] Arioglu-Inan E, Kaykı-Mutlu G, Michel MC. Cardiac β_3 -adrenoceptors - a role in human pathophysiology? *Br J Pharmacol*. 2019; 176(14):2482-2495. <https://doi.org/10.1111/bph.14635>
- [43] Erdogan BR, Michel MC, Arioglu-Inan E. Expression and signaling of β -adrenoceptor subtypes in the diabetic heart. *Cells*. 2020;9(12):2548. <https://doi.org/10.3390/cells9122548>
- [44] Dinçer ÜD, Onay A, Ari N, Özçelikay AT, Altan VM. The effects of diabetes on b-adrenoceptor mediated responsiveness of human and rat atria. *Diabetes Res Clin Pract*. 1998; 40(2):113-122. [https://doi.org/10.1016/S0168-8227\(98\)00034-5](https://doi.org/10.1016/S0168-8227(98)00034-5)
- [45] Dinçer ÜD, Bidasee KR, Güner Ş, Tay A, Özçelikay AT, Altan VM. The effect of diabetes on expression of β_1 -, β_2 -, and β_3 -adrenoreceptors in rat hearts. *Diabetes*. 2001;50(2):455-461. <https://doi.org/10.2337/diabetes.50.2.455>
- [46] Jiang C, Carillion A, Na N, De Jong A, Feldman S, Lacorte JM, Bonnefont-Rousselot D, Riou B, Amour J. Modification of the β -Adrenoceptor Stimulation Pathway in Zucker Obese and Obese Diabetic Rat Myocardium. *Crit Care Med*. 2015;43(7):e241-9. <https://doi.org/10.1097/CCM.0000000000000999>
- [47] Oelze M, Kröller-Schön S, Welschof P, Jansen T, Hausding M, Mikhed Y, Stamm P, Mader M, Zinßius E, Agdauletova S, Gottschlich A, Steven S, Schulz E, Bottari SP, Mayoux E, Münzel T, Daiber A. The sodium-glucose co-transporter 2 inhibitor empagliflozin improves diabetes-induced vascular dysfunction in the streptozotocin diabetes rat model by interfering with oxidative stress and glucotoxicity. *PLoS One*. 2014 ;9(11):e112394. <https://doi.org/10.1371/journal.pone.0112394>
- [48] Steven S, Oelze M, Hanf A, Kröller-Schön S, Kashani F, Roohani S, Welschof P, Kopp M, Gödtel-Armbrust U, Xia N, Li H, Schulz E, Lackner KJ, Wojnowski L, Bottari SP, Wenzel P, Mayoux E, Münzel T, Daiber A. The

- SGLT2 inhibitor empagliflozin improves the primary diabetic complications in ZDF rats. *Redox Biol.* 2017;13:370-385. <https://doi.org/10.1016/j.redox.2017.06.009>
- [49] Wentworth JM, Fourlanos S, Colman PG, Harrison LC. A pilot study of the feasibility of empagliflozin in recent-onset type 1 diabetes. *Metabol Open.* 2020; 100021. <https://doi.org/10.1016/j.metop.2020.100021>
- [50] Rosenstock J, Marquard J, Laffel LM, Neubacher D, Kaspers S, Cherney DZ, Zinman B, Skyler JS, George J, Soleymanlou N, Perkins BA. Empagliflozin as Adjunctive to Insulin Therapy in Type 1 Diabetes: The EASE Trials. *Diabetes Care.* 2018 ;41(12):2560-2569. <https://doi.org/10.2337/dc18-1749>
- [51] Garg SK, Henry RR, Banks P, Buse JB, Davies MJ, Fulcher GR, Pozzilli P, Gesty-Palmer D, Lapuerta P, Simó R, Danne T, McGuire DK, Kushner JA, Peters A, Strumph P. Effects of Sotagliflozin Added to Insulin in Patients with Type 1 Diabetes. *N Engl J Med.* 2017;377(24):2337-2348. <https://doi.org/10.1056/NEJMoa1708337>
- [52] Arioglu Inan E, Ellenbroek JH, Michel MC. A systematic review of urinary bladder hypertrophy in experimental diabetes: part I. streptozotocin-induced rat models. *Neurourol Urodyn.* 2018; 37(4):1212-1219. <https://doi.org/10.1002/nau.23490>
- [53] Onay-Besicki A, Guner S, Arioglu E, Ozakca I, Ozcelikay AT, Altan VM. The effects of chronic trimetazidine treatment on mechanical function and fatty acid oxidation in diabetic rat hearts. *Can J Physiol Pharmacol.* 2007;85(5):527-535. <https://doi.org/10.1139/y07-036>
- [54] Hafez G, Gonulalan U, Kosan M, Arioglu E, Ozturk B, Cetinkaya M, Gur S. Acetylsalicylic acid protects erectile function in diabetic rats. *Andrologia.* 2014;46(9):997-1003. <https://doi.org/10.1111/and.12187>
- [55] Gonulalan U, Kosan M, Hafez G, Arioglu E, Akdemir O, Ozturk B, Gur S, Cetinkaya M. The effect of diabetes mellitus on alpha1-adrenergic receptor subtypes in the bladder of rats. *Urology.* 2012;80(4):951 e9-16. <https://doi.org/10.1016/j.urology.2012.06.019>
- [56] Arioglu-Inan E, Ozakca I, Kayki-Mutlu G, Sepici-Dincel A, Altan VM. The role of insulin-thyroid hormone interaction on beta-adrenoceptor-mediated cardiac responses. *Eur J Pharmacol.* 2013; 718(1-3):533-543. <https://doi.org/10.1016/j.ejphar.2013.06.021>
- [57] Ozakca I, Arioglu E, Guner S, Altan VM, Ozcelikay AT. Role of beta-3-adrenoceptor in catecholamine-induced relaxations in gastric fundus from control and diabetic rats. *Pharmacology.* 2007;80(4):227-238. <https://doi.org/10.1159/000104876>
- [58] Kayki-Mutlu G, Karaomerlioglu I, Arioglu-Inan E. The effect of nitric oxide synthase on beta 3 adrenoceptor mediated relaxation in STZ diabetic rat heart. *J Fac Pharm Ankara.* 2019; 43(1):64-71. <https://doi.org/10.33483/jfpau.519948>
- [59] Kayki-Mutlu G, Arioglu-Inan E, Ozakca I, Ozcelikay AT, Altan VM. beta3-Adrenoceptor-mediated responses in diabetic rat heart. *Gen Physiol Biophys.* 2014; 33(1):99-109. <https://doi.org/10.4149/gpb.2013065>
- [60] Mostafavinia A, Amini A, Ghorishi SK, Pouriran R, Bayat M. The effects of dosage and the routes of administrations of streptozotocin and alloxan on induction rate of type1 diabetes mellitus and mortality rate in rats. *Lab Anim Res.* 2016; 32(3):160-165. <https://doi.org/10.5625/lar.2016.32.3.160>
- [61] Zannad F, Ferreira JP, Pocock SJ, Anker SD, Butler J, Filippatos G, et al. SGLT2 inhibitors in patients with heart failure with reduced ejection fraction: a meta-analysis of the EMPEROR-Reduced and DAPA-HF trials. *Lancet.* 2020;396(10254):819-829. [https://doi.org/10.1016/s0140-6736\(20\)31824-9](https://doi.org/10.1016/s0140-6736(20)31824-9)
- [62] Yesilyurt ZE, Erdogan BR, Karaomerlioglu I, Muderrisoglu AE, Michel MC, Arioglu-Inan E. Urinary Bladder Weight and Function in a Rat Model of Mild Hyperglycemia and Its Treatment With Dapagliflozin. *Front Pharmacol.* 2019; 10:911. <https://doi.org/10.3389/fphar.2019.00911>
- [63] Yesilyurt ZE, Ertürk BM, Erdogan BR, Arioglu Inan E, Michel MC. Effects of the sodium-glucose transporter 2 inhibitor empagliflozin on bladder sie, contraction and ralexation in a rat model of type 1 diabetes. *Neurourol Urodyn.* 2021; 42 Suppl 2:S141-S2. <https://doi.org/10.1002/nau.24746>
- [64] Michel MC, Yesilyurt ZE, Öztürk G, Arioglu Inan E. Empagliflozin and linagliptin do not affect urinary bladder enlargement in a rat model of type 1 diabetes. *Br J Pharmacol.* 2021; 178:4977-4978. <https://doi.org/10.1111/bph.15648>
- [65] Kim DH, Kim YJ, Kim HK, Chang SA, Kim MS, Sohn DW, Oh BH, Park YB. Usefulness of mitral annulus velocity for the early detection of left ventricular dysfunction in a rat model of diabetic cardiomyopathy. *J Cardiovasc Ultrasound.* 2010;18(1):6-11. <https://doi.org/10.4250/jcu.2010.18.1.6>
- [66] Gauthier C, Tavernier G, Charpentier F, Langin D, Le Marec H. Functional beta3-adrenoceptor in the human heart. *J Clin Invest.* 1996; 98(2):556-562. <https://doi.org/10.1172/JCI118823>
- [67] Michel MC, Murphy TJ, Motulsky HJ. New Author Guidelines for Displaying Data and Reporting Data Analysis and Statistical Methods in Experimental Biology. *Mol Pharmacol.* 2020; 97(1):49-60. <https://doi.org/10.1124/mol.119.118927>
- [68] Erdogan BR, Vollert J, Michel MC. Choice of y-axis can mislead readers. *Naunyn Schmiedebergs Arch Pharmacol.* 2020; 393(9):1769-1772. <https://doi.org/10.1007/s00210-020-01926-x>
- [69] Amrhein V, Greenland S, McShane B. Scientists rise up against statistical significance. *Nature.* 2019;567(7748):305-307. <https://doi.org/10.1038/d41586-019-00857-9>
- [70] Michel MC, Murphy TJ, Motulsky HJ. New Author Guidelines for Displaying Data and Reporting Data Analysis and Statistical Methods in Experimental Biology. *J Pharmacol Exp Ther.* 2020; 372(1):136-147. <https://doi.org/10.1124/jpet.119.264143>