Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci.
Mathematics, **44** (1), 41-50 (2024).
https://doi.org/10.30546/2617-7900.44.1.2024.41

Boundedness of the anisotropic fractional maximal operator in total anisotropic Morrey spaces

Vagif S. Guliyev***, Fatai A. Isayev, Ayhan Serbetci**

Received: 21.08.2023 / Revised: 19.03.2024 / Accepted: 06.04.2024

Abstract. *We give necessary and sufficient conditions for the boundedness of the anisotropic fractional maximal operator M_{α}^{d} in total anisotropic Morrey spaces $L_{p,\lambda,\mu}^{d}(\mathbb{R}^n)$.*

Keywords. Total anisotropic Morrey spaces, anisotropic fractional maximal function.

Mathematics Subject Classification (2010): 42B20, 42B25, 42B35.

1 Introduction

Let $\mathbb{R}^n$ be the n-dimension Euclidean space with the norm $|x|$ for each $x \in \mathbb{R}^n$, S^{n-1} denotes the unit sphere on $\mathbb{R}^n$. For $x \in \mathbb{R}^n$ and $r > 0$, let $B(x,r)$ denote the open ball centered at x of radius r and ${}^{\complement}B(x,r)$ denote the set $\mathbb{R}^n \backslash B(x,r)$. Let $d = (d_1, \ldots, d_n)$, $d_i \geq 1$, $i = 1, \ldots, n$, $|d| = \sum_{i=1}^{n} d_i$ and $t^d x \equiv \left(t^{d_1} x_1, \ldots, t^{d_n} x_n\right)$. By [4,6], the function $F(x,\rho) = \sum_{i=1}^{n} x_i^2 \rho^{-2d_i}$, considered for any fixed $x \in \mathbb{R}^n$, is a decreasing one with respect to $\rho > 0$ and the equation $F(x,\rho) = 1$ is uniquely solvable. This unique solution will be denoted by $\rho(x)$. It is a simple matter to check that $\rho(x-y)$ defines a distance between any two points $x,\ y \in \mathbb{R}^n$. Thus $\mathbb{R}^n$, endowed with the metric ρ, defines a homogeneous metric space ([4–6]). The balls with respect to ρ, centered at x of radius r, are just the ellipsoids

$$\mathcal{E}(x,r) \equiv \mathcal{E}_d(x,r) = \left\{ y \in \mathbb{R}^n : \frac{(y_1 - x_1)^2}{r^{2d_1}} + \cdots + \frac{(y_n - x_n)^2}{r^{2d_n}} < 1 \right\},$$

with the Lebesgue measure $|\mathcal{E}(x,r)| = v_n r^{|d|}$, where v_n is the volume of the unit ball in $\mathbb{R}^n$. Let also $\mathit{\Pi}(x,r) \equiv \mathit{\Pi}_d(x,r) = \{y \in \mathbb{R}^n : \max_{1 \leq i \leq n} |x_i - y_i|^{1/d_i} < r\}$ denote

* Corresponding author

V.S. Guliyev
Institute of Applied Mathematics, Baku State University, Baku, Azerbaijan
Department of Mathematics, Ahi Evran University, Kirsehir, Turkey
Institute of Mathematics and Mechanics, Baku, Azerbaijan
E-mail: vagif@guliyev.com

F.A. Isayev
Institute of Mathematics and Mechanics, Baku, Azerbaijan
E-mail: isayevfatai@yahoo.com

A. Serbetci
Department of Mathematics, Ankara University, Ankara, Turkey
E-mail: serbetci@ankara.edu.tr

the parallelopiped, ${}^{\complement}\mathcal{E}(x,r) = \mathbb{R}^n \setminus \mathcal{E}(x,r)$ be the complement of $\mathcal{E}(0,r)$. If $d = \mathbf{1} \equiv (1,\dots,1)$, then clearly $\rho(x) = |x|$ and $\mathcal{E}_{\mathbf{1}}(x,r) = B(x,r)$. Note that in the standard parabolic case $d = (1,\dots,1,2)$ we have

$$\rho(x) = \sqrt{\frac{|x'|^2 + \sqrt{|x'|^4 + x_n^2}}{2}}, \quad x = (x', x_n).$$

Let $f \in L_1^{\mathrm{loc}}(\mathbb{R}^n)$. The anisotropic fractional maximal operator M_α^d is given by

$$M_\alpha^d f(x) = \sup_{t>0} |\mathcal{E}(x,t)|^{-1+\frac{\alpha}{|d|}} \int_{\mathcal{E}(x,t)} |f(y)|dy, \;\; 0 \le \alpha < |d|,$$

where $|\mathcal{E}(x,t)|$ is the Lebesgue measure of the ellipsoid $\mathcal{E}(x,t)$. If $\alpha = 0$, then $M^d \equiv M_0^d$ is the anisotropic Hardy-Littlewood maximal operator. If $d = \mathbf{1}$, then $M_\alpha \equiv M_\alpha^d$ is the fractional maximal operator and $M \equiv M^d$ is the classical Hardy-Littlewood maximal operator.

Morrey spaces, introduced by C. B. Morrey [12], play important roles in the regularity theory of PDE, including heat equations and Navier-Stokes equations. In [10] Guliyev introduce a variant of Morrey spaces called total Morrey spaces $L_{p,\lambda,\mu}(\mathbb{R}^n)$, $0 < p < \infty$, $\lambda \in \mathbb{R}$ and $\mu \in \mathbb{R}$. In [1] Abasova and Omarova consider the total anisotropic Morrey spaces $L_{p,\lambda,\mu}^d(\mathbb{R}^n)$, give basic properties of the spaces $L_{p,\lambda,\mu}^d(\mathbb{R}^n)$ and study some embeddings into the Morrey space $L_{p,\lambda,\mu}^d(\mathbb{R}^n)$. In [11] Guliyev find necessary and sufficient conditions for the boundedness of the fractional maximal operator M_α in the total Morrey spaces $L_{p,\lambda,\mu}(\mathbb{R}^n)$.

The aim of this paper is to give necessary and sufficient conditions for the boundedness of the anisotropic fractional maximal operator M_α^d on total anisotropic Morrey spaces $L_{p,\lambda,\mu}^d(\mathbb{R}^n)$. We prove the strong and weak type Spanne and Adams type boundedness of M_α^d on $L_{p,\lambda,\mu}^d(\mathbb{R}^n)$, respectively.

By $A \lesssim B$ we mean that $A \le CB$ with some positive constant C independent of appropriate quantities. If $A \lesssim B$ and $B \lesssim A$, we write $A \approx B$ and say that A and B are equivalent.

2 Anisotropic fractional maximal operator in total anisotropic Morrey spaces

In this section we find necessary and sufficient conditions for the boundedness of the anisotropic fractional maximal operator M_α^d in the total anisotropic Morrey spaces $L_{p,\lambda,\mu}^d(\mathbb{R}^n)$.

Definition 2.1 *Let $d = (d_1,\dots,d_n)$, $d_i \ge 1$, $i = 1,\dots,n$. Let also $0 < p < \infty$, $\lambda \in \mathbb{R}$, $\mu \in \mathbb{R}$, $[t]_1 = \min\{1,t\}$, $t > 0$. We denote by $L_{p,\lambda}^d(\mathbb{R}^n)$ the anisotropic Morrey space, by $\widetilde{L}_{p,\lambda}^d(\mathbb{R}^n)$ the modified anisotropic Morrey space [9], and by $L_{p,\lambda,\mu}^d(\mathbb{R}^n)$ the total anisotropic Morrey space [1] the set of all classes of locally integrable functions f with the finite norms*

$$\|f\|_{L_{p,\lambda}^d} = \sup_{x\in\mathbb{R}^n,\, t>0} t^{-\frac{\lambda}{p}} \|f\|_{L_p(\mathcal{E}(x,t))},$$

$$\|f\|_{\widetilde{L}_{p,\lambda}^d} = \sup_{x\in\mathbb{R}^n,\, t>0} [t]_1^{-\frac{\lambda}{p}} \|f\|_{L_p(\mathcal{E}(x,t))}$$

and

$$\|f\|_{L^d_{p,\lambda,\mu}} = \sup_{x\in\mathbb{R}^n,\, t>0} [t]_1^{-\frac{\lambda}{p}}\, [1/t]_1^{\frac{\mu}{p}}\, \|f\|_{L_p(\mathcal{E}(x,t))},$$

respectively.

Definition 2.2 *Let $d=(d_1,\dots,d_n)$, $d_i\geq 1$, $i=1,\dots,n$. Let also $0<p<\infty$, $\lambda\in\mathbb{R}$ and $\mu\in\mathbb{R}$. We define the weak anisotropic Morrey space $WL^d_{p,\lambda}(\mathbb{R}^n)$, the weak modified anisotropic Morrey space $W\widetilde{L}^d_{p,\lambda}(\mathbb{R}^n)$ [9] and the weak total anisotropic Morrey space $WL^d_{p,\lambda,\mu}(\mathbb{R}^n)$ [1] as the set of all locally integrable functions f with finite norms*

$$\|f\|_{WL^d_{p,\lambda}} = \sup_{x\in\mathbb{R}^n,\, t>0} t^{-\frac{\lambda}{p}}\, \|f\|_{WL_p(\mathcal{E}(x,t))},$$

$$\|f\|_{W\widetilde{L}^d_{p,\lambda}} = \sup_{x\in\mathbb{R}^n,\, t>0} [t]_1^{-\frac{\lambda}{p}}\, \|f\|_{WL_p(\mathcal{E}(x,t))}$$

and

$$\|f\|_{WL^d_{p,\lambda,\mu}} = \sup_{x\in\mathbb{R}^n,\, t>0} [t]_1^{-\frac{\lambda}{p}}\, [1/t]_1^{\frac{\mu}{p}}\, \|f\|_{WL_p(\mathcal{E}(x,t))},$$

respectively.

Lemma 2.1 *If $0<p<\infty$, $0\leq\lambda\leq|d|$ and $0\leq\mu\leq|d|$, then*

$$L^d_{p,\lambda}(\mathbb{R}^n)\cap L^d_{p,\mu}(\mathbb{R}^n)\subset_{\succ} L^d_{p,\lambda,\mu}(\mathbb{R}^n)$$

and

$$\|f\|_{L^d_{p,\mu,\lambda}} \leq \max\left\{\|f\|_{L^d_{p,\lambda}}, \|f\|_{L^d_{p,\mu}}\right\}.$$

Proof. Let $f\in L^d_{p,\lambda}(\mathbb{R}^n)\cap L^d_{p,\mu}(\mathbb{R}^n)$. Then

$$\begin{aligned}
&\|f\|_{L^d_{p,\lambda,\mu}} = \sup_{x\in\mathbb{R}^n,\, t>0} [t]_1^{-\frac{\lambda}{p}}\, [1/t]_1^{\frac{\mu}{p}}\, \|f\|_{L_p(\mathcal{E}(x,t))}\\
&\max\left\{\sup_{x\in\mathbb{R}^n,0<t\leq 1} t^{-\frac{\lambda}{p}}\, \|f\|_{L_p(\mathcal{E}(x,t))}, \sup_{x\in\mathbb{R}^n,t>1} t^{-\frac{\mu}{p}}\, \|f\|_{L_p(\mathcal{E}(x,t))}\right\}\\
&\leq \max\left\{\|f\|_{L^d_{p,\lambda}}, \|f\|_{L^d_{p,\mu}}\right\}.
\end{aligned}$$

Therefore, $f\in L^d_{p,\lambda,\mu}(\mathbb{R}^n)$ and the embedding $L^d_{p,\lambda}(\mathbb{R}^n)\cap L^d_{p,\mu}(\mathbb{R}^n)\subset_{\succ} L^d_{p,\lambda,\mu}(\mathbb{R}^n)$ is valid.

Lemma 2.2 *If $0<p<\infty$, $0\leq\lambda\leq|d|$ and $0\leq\mu\leq|d|$, then*

$$WL^d_{p,\lambda}(\mathbb{R}^n)\cap WL^d_{p,\mu}(\mathbb{R}^n)\subset_{\succ} WL^d_{p,\lambda,\mu}(\mathbb{R}^n)$$

and

$$\|f\|_{WL^d_{p,\lambda,\mu}(\mathbb{R}^n)} \leq \max\left\{\|f\|_{WL^d_{p,\lambda}}, \|f\|_{WL^d_{p,\mu}}\right\}.$$

The following local estimate is valid.

Lemma 2.3 *[8, Lemma 4.1] Let* $0 \le \alpha < |d|$, $1 \le p < \frac{|d|}{\alpha}$, *and* $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{|d|}$. *Then, for* $p > 1$ *the inequality*

$$\|M_\alpha^d f\|_{L_q(\mathcal{E}(x,r))} \lesssim r^{\frac{|d|}{q}} \sup_{t>2r} t^{-\frac{|d|}{q}} \|f\|_{L_p(\mathcal{E}(x,t))} \tag{2.1}$$

holds for all $\mathcal{E}(x,r)$ *and for all* $f \in L_p^{\mathrm{loc}}(\mathbb{R}^n)$.

Moreover if $p = 1$, *then the inequality*

$$\|M_\alpha^d f\|_{WL_q(\mathcal{E}(x,r))} \lesssim r^{\frac{|d|}{q}} \sup_{t>2r} t^{-\frac{|d|}{q}} \|f\|_{L_1(\mathcal{E}(x,t))} \tag{2.2}$$

holds for all $\mathcal{E}(x,r)$ *and for all* $f \in L_1^{\mathrm{loc}}(\mathbb{R}^n)$.

The following is Spanne's type result for the anisotropic fractional maximal operators in total anisotropic Morrey spaces.

Theorem 2.1 *(Spanne type result) Let* $1 \le p < \infty$, $0 \le \lambda, \mu < |d|$, $0 \le \alpha < \frac{|d|}{p}$ *and* $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{|d|}$.

1. If $p > 1$, $f \in L^d_{p,\lambda,\mu}(\mathbb{R}^n)$, *then* $M_\alpha^d f \in L^d_{q,\frac{\lambda q}{p},\frac{\mu q}{p}}(\mathbb{R}^n)$ *and*

$$\|M_\alpha^d f\|_{L^d_{q,\frac{\lambda q}{p},\frac{\mu q}{p}}} \le C_{p,\lambda,\mu,d} \|f\|_{L^d_{p,\lambda,\mu}}, \tag{2.3}$$

where $C_{p,\lambda,\mu,d}$ *depends only on* p,λ,μ *and* n.

2. If $p = 1$, $f \in L^d_{1,\lambda,\mu}(\mathbb{R}^n)$, *then* $M_\alpha^d f \in WL^d_{q,\lambda q,\mu q}(\mathbb{R}^n)$ *and*

$$\|M_\alpha^d f\|_{WL^d_{q,\lambda q,\mu q}} \le C_{1,\lambda,\mu,d} \|f\|_{L^d_{1,\lambda,\mu}}, \tag{2.4}$$

where $C_{1,\lambda,\mu,d}$ *is independent of* f.

Proof. Let $1 < p < \infty$. From the inequality (2.1) (see Lemma 2.3) we get

$$\begin{aligned}
\|M_\alpha^d f\|_{L^d_{q,\frac{\lambda q}{p},\frac{\mu q}{p}}} &= \sup_{x\in\mathbb{R}^n,\, r>0} [r]_1^{-\frac{\lambda}{p}} [1/r]_1^{\frac{\mu}{p}} \|M_\alpha^d f\|_{L_q(\mathcal{E}(x,r))} \\
&\lesssim \sup_{x\in\mathbb{R}^n,\, r>0} [r]_1^{-\frac{\lambda}{p}} [1/r]_1^{\frac{\mu}{p}} r^{\frac{|d|}{q}} \sup_{t>2r} t^{-\frac{|d|}{q}} \|f\|_{L_p(\mathcal{E}(x,t))} \\
&\lesssim \|f\|_{L^d_{p,\lambda,\mu}} \sup_{r>0} [r]_1^{-\frac{\lambda}{p}} [1/r]_1^{\frac{\mu}{p}} r^{-\alpha+\frac{|d|}{p}} \sup_{t>r} t^{\alpha-\frac{|d|}{p}} [t]_1^{\frac{\lambda}{p}} [1/t]_1^{-\frac{\mu}{p}} \\
&= \|f\|_{L^d_{p,\lambda,\mu}} \sup_{r>0} [r]_1^{-\alpha+\frac{|d|-\lambda}{p}} [1/r]_1^{\alpha-\frac{|d|-\mu}{p}} \sup_{t>r} [t]_1^{\alpha-\frac{|d|-\lambda}{p}} [1/t]_1^{-\alpha+\frac{|d|-\mu}{p}} \\
&= \|f\|_{L^d_{p,\lambda,\mu}},
\end{aligned}$$

which implies that the operator M_α^d is bounded from $L^d_{p,\lambda,\mu}(\mathbb{R}^n)$ to $L^d_{q,\frac{\lambda q}{p},\frac{\mu q}{p}}(\mathbb{R}^n)$.

Let $p=1$. From the inequality (2.2) (see Lemma 2.3) we get

$$\begin{aligned}
&\|M_\alpha^d f\|_{WL^d_{q,\lambda q,\mu q}} = \sup_{x\in\mathbb{R}^n,\, r>0} [r]_1^{-\lambda}\,[1/r]_1^{\mu}\,\|M_\alpha^d f\|_{WL_q(\mathcal{E}(x,r))}\\
&\lesssim \sup_{x\in\mathbb{R}^n,\, r>0} [r]_1^{-\lambda}\,[1/r]_1^{\mu}\, r^{\frac{|d|}{q}} \sup_{t>2r} t^{-\frac{|d|}{q}}\,\|f\|_{L_1(\mathcal{E}(x,t))}\\
&\lesssim \|f\|_{L^d_{1,\lambda,\mu}} \sup_{r>0}[r]_1^{-\lambda}\,[1/r]_1^{\mu}\, r^{-\alpha+n} \sup_{t>r} t^{\alpha-|d|}\,[t]_1^{\lambda}\,[1/t]_1^{-\mu}\\
&= \|f\|_{L^d_{1,\lambda,\mu}} \sup_{r>0}[r]_1^{-\alpha+|d|-\lambda}\,[1/r]_1^{\alpha-(|d|-\mu)} \sup_{t>r}[t]_1^{\alpha-(|d|-\lambda)}\,[1/t]_1^{-\alpha+(|d|-\mu)}\\
&= \|f\|_{L^d_{1,\lambda,\mu}},
\end{aligned}$$

which implies that the operator M_α^d is bounded from $L^d_{1,\lambda,\mu}(\mathbb{R}^n)$ to $WL^d_{q,\lambda q,\mu q}(\mathbb{R}^n)$.

From Theorem 2.1 in the case $\alpha=0$ we get the following corollaries.

Corollary 2.1 *[10, Theorem 1] Let $1\le p<\infty$, $0\le\lambda<|d|$ and $0\le\mu<|d|$.*
1. If $p>1$, $f\in L^d_{p,\lambda,\mu}(\mathbb{R}^n)$, then $M^d f\in L^d_{p,\lambda,\mu}(\mathbb{R}^n)$ and

$$\|M^d f\|_{L^d_{p,\lambda,\mu}} \le C_{p,\lambda,\mu,d}\,\|f\|_{L^d_{p,\lambda,\mu}},$$

where $C_{p,\lambda,\mu,d}$ depends only on p, λ, μ, d and n.
2. If $f\in L^d_{1,\lambda,\mu}(\mathbb{R}^n)$, then $M^d f\in WL^d_{1,\lambda,\mu}(\mathbb{R}^n)$ and

$$\|M^d f\|_{WL^d_{1,\lambda,\mu}} \le C_{1,\lambda,\mu,d}\,\|f\|_{L^d_{1,\lambda,\mu}},$$

where $C_{1,\lambda,\mu,d}$ depends only on p, λ, μ, d and n.

From Theorem 2.1 in the case $\lambda=\mu$ or $\mu=0$ we get the following corollaries.

Corollary 2.2 *[3] Let $1\le p<\infty$, $0\le\lambda<|d|$, $0\le\alpha<\frac{|d|}{p}$ and $\frac{1}{p}-\frac{1}{q}=\frac{\alpha}{|d|}$.*
1. If $p>1$, $f\in L^d_{p,\lambda}(\mathbb{R}^n)$, then $M^d_\alpha f\in L^d_{q,\frac{\lambda q}{p}}(\mathbb{R}^n)$ and

$$\|M^d_\alpha f\|_{L^d_{q,\frac{\lambda q}{p}}} \le C_{p,\lambda,d}\,\|f\|_{L^d_{p,\lambda}}, \tag{2.5}$$

where $C_{p,\lambda,d}$ depends only on p, λ, d and n.
2. If $p=1$, $f\in L^d_{1,\lambda}(\mathbb{R}^n)$, then $Mf\in WL^d_{q,\lambda}(\mathbb{R}^n)$ and

$$\|M^d_\alpha f\|_{WL^d_{q,\lambda q}} \le C_{1,\lambda,d}\,\|f\|_{L^d_{1,\lambda}}, \tag{2.6}$$

where $C_{1,\lambda,d}$ is independent of f.

Corollary 2.3 *[9, Theorem 2.1] Let $1\le p<\infty$, $0\le\lambda<|d|$, $0\le\alpha<\frac{|d|}{p}$ and $\frac{1}{p}-\frac{1}{q}=\frac{\alpha}{|d|}$.*
1. If $p>1$, $f\in \widetilde{L}^d_{p,\lambda}(\mathbb{R}^n)$, then $M^d_\alpha f\in \widetilde{L}^d_{q,\frac{\lambda q}{p}}(\mathbb{R}^n)$ and

$$\|M^d_\alpha f\|_{\widetilde{L}^d_{q,\frac{\lambda q}{p}}} \le C_{p,\lambda,d}\,\|f\|_{\widetilde{L}^d_{p,\lambda}}, \tag{2.7}$$

where $C_{p,\lambda,d}$ depends only on p, λ and n.

2. If $p=1$, $f\in \widetilde{L}^d_{1,\lambda}(\mathbb{R}^n)$, *then* $M^d_\alpha f\in W\widetilde{L}^d_{q,\lambda}(\mathbb{R}^n)$ *and*

$$\|M^d_\alpha f\|_{W\widetilde{L}^d_{q,\lambda_q}}\le C_{1,\lambda,d}\,\|f\|_{\widetilde{L}^d_{1,\lambda}}, \tag{2.8}$$

where $C_{1,\lambda,d}$ *is independent of* f.

The following is Adam's type result for the anisotropic fractional maximal operators in total anisotropic Morrey spaces.

Theorem 2.2 *(Adams type result) Let* $1\le p<\infty$, $0\le\lambda\le\mu<|d|$, $0\le\alpha<\frac{|d|-\mu}{p}$.

1) If $1<p<\frac{|d|-\mu}{\alpha}$, *then condition* $\frac{\alpha}{|d|-\lambda}\le\frac{1}{p}-\frac{1}{q}\le\frac{\alpha}{|d|-\mu}$ *is necessary and sufficient for the boundedness of the operator* M^d_α *from* $L^d_{p,\lambda,\mu}(\mathbb{R}^n)$ *to* $L^d_{q,\lambda,\mu}(\mathbb{R}^n)$.

2) If $p=1<\frac{|d|-\mu}{\alpha}$, *then condition* $\frac{\alpha}{|d|-\lambda}\le 1-\frac{1}{q}\le\frac{\alpha}{|d|-\mu}$ *is necessary and sufficient for the boundedness of the operator* M^d_α *from* $L^d_{1,\lambda,\mu}(\mathbb{R}^n)$ *to* $WL^d_{q,\lambda,\mu}(\mathbb{R}^n)$.

Proof. *Sufficiency.* Let $1\le p<\frac{|d|-\mu}{\alpha}$, $\frac{\alpha}{|d|-\lambda}\le\frac{1}{p}-\frac{1}{q}\le\frac{\alpha}{|d|-\mu}$ and $f\in L^d_{p,\lambda,\mu}(\mathbb{R}^n)$.

$$\begin{aligned}
M^d_\alpha f(x)&\approx\sup_{r>0} r^{\alpha-|d|}\,\|f\|_{L_1(\mathcal{E}(x,r))}\\
&\le\sup_{r>0}\min\{r^\alpha\,M^d f(x),\,r^{\alpha-\frac{|d|}{p}}\,\|f\|_{L_p(\mathcal{E}(x,r))}\}\\
&\le\sup_{r>0}\min\{r^\alpha\,M^d f(x),\,r^{\alpha-\frac{|d|}{p}}\,[r]_1^{\frac{\lambda}{p}}\,[1/r]_1^{-\frac{\mu}{p}}\,\|f\|_{L^d_{p,\lambda,\mu}}\}\\
&\le\sup_{r>0}\min\{r^\alpha\,M^d f(x),\,[r]_1^{\alpha-\frac{|d|-\lambda}{p}}\,[1/r]_1^{-\alpha+\frac{|d|-\mu}{p}}\,\|f\|_{L^d_{p,\lambda,\mu}}\}\\
&\le\max\Big\{\sup_{0<r\le1}\min\{r^\alpha\,M^d f(x),\,r^{\alpha-\frac{|d|-\lambda}{p}}\,\|f\|_{L^d_{p,\lambda,\mu}}\},\\
&\qquad\sup_{r>1}\min\{r^\alpha\,M^d f(x),\,r^{\alpha-\frac{|d|-\mu}{p}}\,\|f\|_{L^d_{p,\lambda,\mu}}\}\Big\}.
\end{aligned}$$

Minimizing with respect to r, at

$$r=\Big(\frac{\|f\|_{L^d_{p,\lambda,\mu}}}{M^d f(x)}\Big)^{\frac{p}{|d|-\lambda}}\quad\text{and}\quad r=\Big(\frac{\|f\|_{L^d_{p,\lambda,\mu}}}{M^d f(x)}\Big)^{\frac{p}{|d|-\mu}}$$

we have

$$\begin{aligned}
M^d_\alpha f(x)\le\max\Big\{&\big(M^d f(x)\big)^{1-\frac{\alpha p}{|d|-\lambda}}\,\|f\|^{\frac{\alpha p}{|d|-\lambda}}_{L^d_{p,\lambda,\mu}},\\
&\big(Mf(x)\big)^{1-\frac{\alpha p}{|d|-\mu}}\,\|f\|^{\frac{\alpha p}{|d|-\mu}}_{L^d_{p,\lambda,\mu}}\Big\},
\end{aligned}\tag{2.9}$$

where we have used that the supremum is achieved when the minimum parts are balanced. From Corollary 2.1 and inequality (2.9), we get

$$\begin{aligned}
\|M^d_\alpha f\|_{L^d_{q,\lambda,\mu}}&\lesssim\|f\|^{1-\frac{p}{q}}_{L^d_{p,\lambda,\mu}}\,\|(M^d f)^{\frac{p}{q}}\|_{L^d_{q,\lambda,\mu}}\\
&=\|f\|^{1-\frac{p}{q}}_{L_{p,\lambda,\mu}}\,\|M^d f\|^{\frac{p}{q}}_{L^d_{p,\lambda,\mu}}\lesssim\|f\|_{L^d_{p,\lambda,\mu}},
\end{aligned}$$

if $1 < p < q < \infty$ and

$$\|M_\alpha^d f\|_{WL^d_{q,\lambda,\mu}} \lesssim \|f\|^{1-\frac{1}{q}}_{L^d_{1,\lambda,\mu}} \|M^d f\|^{\frac{1}{q}}_{WL^d_{1,\lambda,\mu}} \lesssim \|f\|_{L^d_{1,\lambda,\mu}},$$

if $p = 1 < q < \infty$.

Necessity. Let $1 < p < \frac{|d|-\mu}{\alpha}$, $\frac{\alpha}{|d|-\lambda} \le \frac{1}{p} - \frac{1}{q} \le \frac{\alpha}{|d|-\mu}$, $f \in L^d_{p,\lambda,\mu}(\mathbb{R}^n)$ and assume that M^d_α is bounded from $L^d_{p,\lambda,\mu}(\mathbb{R}^n)$ to $L^d_{q,\lambda,\mu}(\mathbb{R}^n)$.

Define $f_{t^d}(x) =: f(t^d x)$, $[t]_{1,+} = \max\{1, t\}$. Then

$$\left\|f_{t^d}\right\|_{L^d_{p,\lambda,\mu}} = \sup_{x\in\mathbb{R}^n, r>0} [r]_1^{-\frac{\lambda}{p}} [1/r]_1^{\frac{\mu}{p}} \|f_{t^d}\|_{L_p(\mathcal{E}(x,r))}$$

$$= t^{-\frac{|d|}{p}} \sup_{x\in\mathbb{R}^n, r>0} [r]_1^{-\frac{\lambda}{p}} [1/r]_1^{\frac{\mu}{p}} \|f\|_{L_p(\mathcal{E}(x,tr))}$$

$$= t^{-\frac{|d|}{p}} \sup_{r>0} \Big(\frac{[tr]_1}{[r]_1}\Big)^{\frac{\lambda}{p}} \sup_{r>0} \Big(\frac{[1/r]_1}{[1/(tr)]_1}\Big)^{\frac{\mu}{p}}$$

$$\times \sup_{x\in\mathbb{R}^n, r>0} [tr]_1^{-\frac{\lambda}{p}} [1/(tr)]_1^{\frac{\mu}{p}} \|f\|_{L_p(\mathcal{E}(x,tr))}$$

$$= t^{-\frac{|d|}{p}} [t]_{1,+}^{\frac{\lambda}{p}} [1/t]_{1,+}^{-\frac{\mu}{p}} \|f\|_{L^d_{p,\lambda,\mu}},$$

and

$$M^d_\alpha f_{t^d}(x) = t^{-\alpha} M^d_\alpha f(t^d x),$$

$$\left\|M^d_\alpha f_{t^d}\right\|_{L^d_{q,\lambda,\mu}} = t^{-\alpha} \sup_{x\in\mathbb{R}^n, r>0} [r]_1^{-\frac{\lambda}{p}} [1/r]_1^{\frac{\mu}{p}} \|M^d_\alpha f(t^d\,\cdot)\|_{L_q(\mathcal{E}(x,r))}$$

$$= t^{-\alpha-\frac{|d|}{q}} \sup_{r>0} \Big(\frac{[tr]_1}{[r]_1}\Big)^{\lambda/q} \sup_{r>0} \Big(\frac{[1/r]_1}{[1/(tr)]_1}\Big)^{\mu/q}$$

$$\times \sup_{x\in\mathbb{R}^n, r>0} [tr]_1^{-\frac{\lambda}{p}} [1/(tr)]_1^{\frac{\mu}{p}} \|M^d_\alpha f\|_{L_q(\mathcal{E}(t^d x,tr))}$$

$$= t^{-\alpha-\frac{|d|}{q}} [t]_{1,+}^{\frac{\lambda}{q}} [1/t]_{1,+}^{-\frac{\mu}{q}} \left\|M^d_\alpha f\right\|_{L^d_{q,\lambda,\mu}}.$$

By the boundedness of M^d_α from $L^d_{p,\lambda,\mu}(\mathbb{R}^n)$ to $L^d_{q,\lambda,\mu}(\mathbb{R}^n)$ we have

$$\left\|M^d_\alpha f\right\|_{L^d_{q,\lambda,\mu}} = t^{\alpha+\frac{|d|}{q}} [t]_{1,+}^{-\frac{\lambda}{q}} [1/t]_{1,+}^{\frac{\mu}{q}} \left\|M^d_\alpha f_{t^d}\right\|_{L^d_{q,\lambda,\mu}}$$

$$\lesssim t^{\alpha+\frac{|d|}{q}}\,[t]_{1,+}^{-\frac{\lambda}{q}}\,[1/t]_{1,+}^{\frac{\mu}{q}}\,\left\|f_{t^d}\right\|_{L_{p,\lambda,\mu}}$$

$$= t^{\alpha+\frac{|d|}{q}-\frac{|d|}{p}}\,[t]_{1,+}^{\frac{\lambda}{p}-\frac{\lambda}{q}}\,[1/t]_{1,+}^{-\frac{\mu}{p}+\frac{\mu}{q}}\,\|f\|_{L_{p,\lambda,\mu}}$$

$$= t^{\alpha}\,[t]_{1,+}^{-\frac{|d|-\lambda}{p}+\frac{|d|-\lambda}{q}}\,[1/t]_{1,+}^{\frac{|d|-\mu}{p}-\frac{|d|-\mu}{q}}\,\|f\|_{L^d_{p,\lambda,\mu}}.$$

If $\frac{1}{p} < \frac{1}{q} + \frac{\alpha}{|d|-\mu}$, then by letting $t \to 0$ we have $\left\|M^d_\alpha f\right\|_{L^d_{q,\lambda,\mu}} = 0$ for all $f \in L^d_{p,\lambda,\mu}(\mathbb{R}^n)$.

As well as if $\frac{1}{p} > \frac{1}{q} + \frac{\alpha}{|d|-\lambda}$, then at $t \to \infty$ we obtain $\left\|M^d_\alpha f\right\|_{L^d_{q,\lambda,\mu}} = 0$ for all $f \in L^d_{p,\lambda,\mu}(\mathbb{R}^n)$.

Therefore $\frac{\alpha}{|d|-\lambda} \le \frac{1}{p} - \frac{1}{q} \le \frac{\alpha}{|d|-\mu}$.

Let $p = 1 < \frac{|d|-\mu}{\alpha}$, $f \in L^d_{p,\lambda,\mu}(\mathbb{R}^n)$ and assume that M^d_α is bounded from $L^d_{1,\lambda,\mu}(\mathbb{R}^n)$ to $WL^d_{q,\lambda,\mu}(\mathbb{R}^n)$. Then

$$\left\|f_{t^d}\right\|_{L^d_{1,\lambda,\mu}} = t^{-|d|}\,[t]_{1,+}^{\lambda}\,[1/t]_{1,+}^{-\mu}\,\|f\|_{L^d_{1,\lambda,\mu}}$$

and

$$\left\|M^d_\alpha f_{t^d}\right\|_{WL^d_{q,\lambda,\mu}} = t^{-\alpha}\sup_{x\in\mathbb{R}^n,\,r>0}[r]_1^{-\frac{\lambda}{p}}\,[1/r]_1^{\frac{\mu}{p}}\,\|M^d_\alpha f(t^d\,\cdot)\|_{WL_q(\mathcal{E}(x,r))}$$

$$= t^{-\alpha-\frac{|d|}{q}}\sup_{r>0}\Big(\frac{[tr]_1}{[r]_1}\Big)^{\lambda/q}\sup_{r>0}\Big(\frac{[1/r]_1}{[1/(tr)]_1}\Big)^{\mu/q}$$

$$\times\sup_{x\in\mathbb{R}^n,\,r>0}[tr]_1^{-\frac{\lambda}{p}}\,[1/(tr)]_1^{\frac{\mu}{p}}\,\|M^d_\alpha f\|_{WL_q(\mathcal{E}(t^d x,tr))}$$

$$= t^{-\alpha-\frac{n}{q}}\,[t]_{1,+}^{\frac{\lambda}{q}}\,[1/t]_{1,+}^{-\frac{\mu}{q}}\,\left\|M^d_\alpha f\right\|_{WL^d_{q,\lambda,\mu}}.$$

By the boundedness of M^d_α from $L^d_{1,\lambda,\mu}(\mathbb{R}^n)$ to $WL^d_{q,\lambda,\mu}(\mathbb{R}^n)$ we have

$$\left\|M^d_\alpha f\right\|_{WL^d_{q,\lambda,\mu}} = t^{\alpha+\frac{|d|}{q}}\,[t]_{1,+}^{-\frac{\lambda}{q}}\,[1/t]_{1,+}^{\frac{\mu}{q}}\,\left\|M^d_\alpha f_{t^d}\right\|_{WL^d_{q,\lambda,\mu}}$$

$$\lesssim t^{\alpha+\frac{|d|}{q}}\,[t]_{1,+}^{-\frac{\lambda}{q}}\,[1/t]_{1,+}^{\frac{\mu}{q}}\,\left\|f_{t^d}\right\|_{L^d_{1,\lambda,\mu}}$$

$$= t^{\alpha+\frac{|d|}{q}-|d|}\,[t]_{1,+}^{\lambda-\frac{\lambda}{q}}\,[1/t]_{1,+}^{-\mu+\frac{\mu}{q}}\,\|f\|_{L^d_{1,\lambda,\mu}}$$

$$= t^{\alpha}\,[t]_{1,+}^{-|d|+\lambda+\frac{|d|-\lambda}{q}}\,[1/t]_{1,+}^{|d|-\mu-\frac{|d|-\mu}{q}}\,\|f\|_{L^d_{1,\lambda,\mu}}.$$

If $1 < \frac{1}{q} + \frac{\alpha}{|d|-\lambda}$, then by letting $t \to 0$ we have $\left\|M_\alpha^d f\right\|_{WL^d_{q,\lambda,\mu}} = 0$ for all $f \in L^d_{1,\lambda,\mu}(\mathbb{R}^n)$.

As well as if $1 > \frac{1}{q} + \frac{\alpha}{|d|-\mu}$, then at $t \to \infty$ we obtain $\left\|M_\alpha^d f\right\|_{WL^d_{q,\lambda,\mu}} = 0$ for all $f \in L^d_{1,\lambda,\mu}(\mathbb{R}^n)$.

Therefore $\frac{\alpha}{|d|-\lambda} \leq 1 - \frac{1}{q} \leq \frac{\alpha}{|d|-\mu}$.

From Theorem 2.2 in the case $\lambda = \mu$ or $\mu = 0$ we get the following corollaries.

Corollary 2.4 *[2, Theorem 3.1] (Adams result)*

Let $1 \leq p < \infty$, $0 \leq \lambda < |d|$, $0 \leq \alpha < \frac{|d|-\lambda}{p}$.

1) If $1 < p < \frac{|d|-\lambda}{\alpha}$, *then the condition* $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{|d|-\lambda}$ *is necessary and sufficient for the boundedness of the operator* M_α^d *from* $L^d_{p,\lambda}(\mathbb{R}^n)$ *to* $L^d_{q,\lambda}(\mathbb{R}^n)$.

2) If $p = 1 < \frac{|d|-\lambda}{\alpha}$, *then the condition* $1 - \frac{1}{q} = \frac{\alpha}{|d|-\lambda}$ *is necessary and sufficient for the boundedness of the operator* M_α^d *from* $L^d_{1,\lambda}(\mathbb{R}^n)$ *to* $WL^d_{q,\lambda}(\mathbb{R}^n)$.

Corollary 2.5 *[7, Corollary 1] Let* $1 \leq p < \infty$, $0 \leq \lambda < |d|$, $0 \leq \alpha < \frac{|d|-\lambda}{p}$.

1) If $1 < p < \frac{|d|-\lambda}{\alpha}$, *then the condition* $\frac{\alpha}{|d|} \leq \frac{1}{p} - \frac{1}{q} \leq \frac{\alpha}{|d|-\lambda}$ *is necessary and sufficient for the boundedness of the operator* M_α^d *from* $\widetilde{L}^d_{p,\lambda}(\mathbb{R}^n)$ *to* $\widetilde{L}^d_{q,\lambda}(\mathbb{R}^n)$.

2) If $p = 1 < \frac{|d|-\lambda}{\alpha}$, *then the condition* $\frac{\alpha}{|d|} \leq 1 - \frac{1}{q} \leq \frac{\alpha}{|d|-\lambda}$ *is necessary and sufficient for the boundedness of the operator* M_α^d *from* $\widetilde{L}^d_{1,\lambda}(\mathbb{R}^n)$ *to* $W\widetilde{L}^d_{q,\lambda}(\mathbb{R}^n)$.

Remark 2.1 Note that in the case of $d = \mathbf{1} \equiv (1, \ldots, 1)$ from Theorem 2.1 we get [11, Theorem 2.1] and from Theorem 2.2 we get [11, Theorem 2.2].

Acknowledgements The authors thank the referee(s) for careful reading the paper and useful comments.

References

1. Abasova, G.A., Omarova, M.N.: *Commutator of anisotropic maximal function with BMO functions on total anisotropic Morrey spaces*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci., Mathematics **43**(1), 3-15 (2022).
2. Adams, D.R.: *A note on Riesz potentials*, Duke Math. J. **42**(4), 765778 (1975).
3. Akbulut, A., Guliyev, V.S., Muradova, Sh.A.: *Boundedness of the anisotropic Riesz potential in anisotropic local Morrey-type spaces*, Complex Var. Elliptic Equ. **58**(2), 259-280 (2013).
4. Besov, O.V., Il'in, V.P., Lizorkin, P.I.: *The L_p-estimates of a certain class of non-isotropically singular integrals,* (Russian) Dokl. Akad. Nauk SSSR, **169**, 1250-1253 (1966).
5. Bramanti, M., Cerutti, M.C.: *Commutators of singular integrals on homogeneous spaces*, Boll. Un. Mat. Ital. B, **10**(7), 843-883 (1996).
6. Fabes, E.B., Rivère, N.: *Singular integrals with mixed homogeneity*, Studia Math. **27**, 19-38 (1966).
7. Guliyev, V.S., Hasanov, J.J., Zeren, Y.: *Necessary and sufficient conditions for the boundedness of the Riesz potential in modified Morrey spaces*, J. Math. Inequal. **5**(4), 491-506 (2011).

8. Guliyev, V.S., Akbulut, A., Mammadov, Y.Y.: *Boundedness of fractional maximal operator and their higher order commutators in generalized Morrey spaces on Carnot groups*, Acta Math. Sci. Ser. B Engl. Ed. **33**(5), 13291346 (2013).
9. Guliyev, V.S., Rahimova, K.R.: *Parabolic fractional maximal operator in modified parabolic Morrey spaces*, J. Funct. Space Appl. Volume 2012, Article ID 543475, 20 pages (2012). doi:10.1155/2012/543475
10. Guliyev, V.S.: *Maximal commutator and commutator of maximal function on total Morrey spaces*, J. Math. Inequal. **16**(4), 15091524 (2022).
11. Guliyev, V.S.: *Characterizations of Lipschitz functions via the commutators of maximal function in total Morrey spaces*, Math. Meth. Appl. Sci. 114 (2024), DOI 10.1002/mma.10038.
12. Morrey, C.B.: *On the solutions of quasi-linear elliptic partial differential equations*, Trans. Amer. Math. Soc. **43**, 126-166 (1938).